

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
17 July 2003 (17.07.2003)

PCT

(10) International Publication Number
WO 03/057214 A1

(51) International Patent Classification⁷: **A61K 31/4166**,
31/404, A61P 35/00, C07D 403/06

(74) Agents: **EFRON, Margaret** et al.; Saliwanchik, Lloyd &
Saliwanchik, Suite A-1, 2421 N.W. 41st Street, Gainesville,
FL 32606-6669 (US).

(21) International Application Number: PCT/US02/35751

(22) International Filing Date:
7 November 2002 (07.11.2002)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/344,564 28 December 2001 (28.12.2001) US
60/344,563 28 December 2001 (28.12.2001) US

(71) Applicant (for all designated States except US): **SOMATOCOR PHARMACEUTICALS, INC.** [US/US]; 8995
Westside Parkway, Ste 200, Alpharetta, GA 30004 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **SHAPIRO, Gideon**
[US/US]; 5507 NW 80th Avenue, Gainesville, FL 32653
(US). **NATCHUS, Michael, G.** [US/US]; 55 Waterview
Trail, Alpharetta, GA 30022 (US). **LOCKWOOD, Mark,**
A. [US/US]; 5315 John's View Street, Alpharetta, GA
30005 (US). **JURCZYCK, Simona** [DE/US]; 4830 N.W.
43rd Street, Gainesville, FL 32606 (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,
MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG,
SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ,
VC, VN, YU, ZA, ZM, ZW.


(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE,
ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK,
TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ,
GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: HYDANTOIN DERIVATIVES WITH AFFINITY FOR SOMATOSTATIN RECEPTORS



(57) Abstract: Non-peptide somatostatin receptor ligands of Formula (I) with conformationally restricted side chains exhibiting high binding affinity toward somatostatin receptors are provided. The compounds exhibit a high selectivity and act as agonists at human subtype 2 somatostatin receptors. The compounds are long acting for advantageous use as medicaments in peripheral diseases where somatostatinergic therapy is indicated. Furthermore, many of the compounds are lipophilic and are particularly useful for treating central nervous system and ophthalmic diseases where penetration of the blood brain and blood retinal barriers is required. It is a further object to describe the preferred stereoisomers of these somatostatin agonists and processes for their preparation. Further objects will become apparent from reading the following description.

Description

HYDANTOIN DERIVATIVES WITH AFFINITY FOR SOMATOSTATIN RECEPTORS

5

Background of the Invention

Somatostatin is an endogenous peptide that performs a variety of important functions within the body. Somatostatin is a highly flexible cyclic peptide with a very short biological half-life. Two known biologically active forms of somatostatin are a 14mer (SRIF-14) and a 28mer N-terminal extended form (SRIF-28). Naturally-occurring
10 somatostatin (also known as somatotropin release inhibiting factor, SRIF) was originally isolated from ovine hypothalamus.

Somatostatin, originally discovered to act as a classical endocrine hormone of the hypothalamic-pituitary system, has since been shown to act additionally as a paracrine and autocrine signaling factor on a wide variety of cell types. The numerous
15 physiological processes currently recognized to be influenced by somatostatin include hormone and peptide factor secretion, neurotransmission, cell proliferation, smooth muscle contraction, nutrient absorption and inflammation. Hormones and peptides regulated by somatostatin include growth hormone (GH), thyroid-stimulating hormone (TSH), prolactin (PRL), insulin, and substance P (SP).

Somatostatin affects the function of many important biological systems such as the endocrine, gastrointestinal, vascular, and immune systems along with the central and peripheral nervous systems. In the endocrine system, somatostatin plays an important role in controlling growth hormone, insulin and glucagon secretion (Koerker *et al.*,
20 *Science* 1974, 184, 482-484). The effects of somatostatin on the gastrointestinal and vascular biological systems have led to clinical applications for somatostatin therapeutics in both of these areas. In the central nervous system (CNS), somatostatin appears to be an important regulator of cognitive functions (Schettini, *Pharmacological Research* 1991, 23, 203-215) and, in specific areas of the brain, appears to act as a neurotransmitter or as a neuromodulator regulating the release of neurotransmitters such as acetylcholine (Gray
25 *et al.*, *J. of Neuroscience* 1990, 10, 2687-2698) and dopamine (Thal *et al.*, *Brain Research* 1986, 372, 205-209). In the peripheral nervous system (PNS), somatostatin is present in catecholamine containing fibers and in sensory terminals together with
30

substance P (Green *et al.*, *Neuroscience* 1992, 50, 745-749) and acts to inhibit their release and mediated effects.

Somatostatin is expressed in diverse cell types of the immune and hematopoietic systems (Hofland *et al.*, *Ann. Medicine* 1999, 31 Suppl. 2, 23-27) and has been shown to
5 inhibit cytokine secretion (*e.g.*, IL-1 β) from peripheral blood mononuclear cells (Peluso
et al., *Neuropeptides* 1996, 30, 443-451) and inhibit the production of immunoglobulin
(Ig)E from B lymphocytes (Kimata *et al.*, *J. Immunology* 1993, 150, 4630-4640).
Furthermore, it has been demonstrated that somatostatin is important as an endogenous
inhibitor of cell proliferation in various normal and neoplastic tissues (Reubi *et al.*,
10 *Trends in Pharmacological Sciences* 1995, 16, 110-115).

The biological effects of somatostatin are mediated through five somatostatin G-
protein receptor subtypes, SSTR1-5 (Reisine *et al.*, *Endocrine Reviews* 1995, 16, 427-
442), that are highly conserved across different species and can be grouped into two
families SSTR 2, 3, and 5, and SSTR 1 and 4. All five receptors are heterogeneously
15 distributed and pharmacologically distinct. Studies utilizing subtype selective
somatostatin receptor agonists have provided evidence that somatostatin subtype 2
receptors (SSTR2) mediate the inhibition of growth hormone release from the anterior
pituitary and of glucagon release from the pancreas; whereas SSTR5 receptors mediate
inhibition of insulin release.

20 Like somatostatin itself, somatostatin receptors have been localized to a wide
variety of tissues and cell types including those belonging to the CNS, PNS, endocrine,
gastrointestinal, vascular, and immune systems. A high incidence of somatostatin
receptors has also been demonstrated in a variety of human tumors. Neuroendocrine
tumors are one class of tumors that exhibit a high density of functionally active
25 somatostatin receptors. Functionally active neuroendocrine tumors present with clinical
symptoms such as gastrinoma and glucagonoma syndrome due to excessive hormone
release from the tumor cell. Such symptoms may be treated through somatostatin
receptor activation.

Another group of tumors recognized as having somatostatin receptors originates
30 in the CNS, which includes both the brain and the spinal cord. While most brain tumors
contain somatostatin receptors, their receptor content often varies depending on tumor
type. Generally, tumors of glial or meningeal origin, including medulloblastomas,
oligodendrogliomas, and differentiated astrocytomas, display somatostatin receptors.

Additional tumors characterized as displaying somatostatin receptors include kidney, breast, and lymphomas, endometrial (Mishima *et al.*, *American Journal of Obstetrics and Gynecology* 1999, 181, 583-590), ovarian (Halmos *et al.*, *Journal of Clinical Endocrinology and Metabolism* 2000, 85, 3509-3512), small cell lung, and prostate (Reubi *et al.*, *Trends in Pharmacological Sciences*, 1995, 16, 110-115).

It has been reported that somatostatin may act directly on cells, *e.g.*, inducing cell death by apoptosis (Srikant, *Biochemical and Biophysical Research Communications* 1995, 209, 400-406). Alternatively, somatostatin may influence tumor growth by indirect mechanisms including effects on growth factor levels that correlate with tumor growth, *e.g.*, IGF-1, or by blocking angiogenesis via inhibiting the proliferation of vascular endothelial cells (Watson *et al.*, *British Journal of Cancer* 2001, 85, 266-272).

Along with somatostatin (SRIF-14) itself, several somatostatin peptide analogs are currently available for various clinical uses. Such uses have been established for cyclic octapeptide therapeutic analogues, including for example, octreotide (Bauer *et al.*, *Life Sciences* 1982, 31,1133), lanreotide (Guisti *et al.*, *European Journal of Clinical Investigation* 1997, 27, 277), and vapreotide (Liebow *et al.*, *Proceedings of the National Academy of Sciences* 1989, 86, 2003). These compounds were originally characterized by their binding effects to prepared rat brain membrane and their effects on endocrine parameters in animals. Structure activity studies on somatostatin peptide analogs have demonstrated that the amino acid side chains from the Phe-Trp-Lys tripeptide subsequence of somatostatin play a key role in receptor binding affinity. The three dimensional structure of octreotide has been determined by x-ray crystallographic studies (see Pohl *et al.*, *Acta Crystallographica* 1995, D51, 48-49). Nuclear magnetic resonance (NMR) structure studies have been performed on other somatostatin peptide analogs (Kessler *et al.*, *Journal of the American Chemical Society* 1983, 105, 6944-6952). These studies indicate that the three dimensional geometry of the amino acid side chains based on the somatostatin Phe-Trp-Lys tripeptide subsequence is defined by a β -turn structure.

Many of the therapeutic properties of somatostatin and somatostatin peptide analogs discussed above can be correlated with the functional activation of the SSSTR2 somatostatin receptor subtype as characterized using *in vitro* cellular constructs which have been transfected with the five individual cloned human somatostatin receptor subtypes.

Current somatostatin peptide therapeutics suffer from numerous drawbacks. Major drawbacks include lack of oral activity, relatively short plasma half life, and poor penetration of the blood retinal barrier (BRB) and of the blood brain barrier (BBB) to access the central nervous system. As a result, somatostatin peptide analog therapeutics are administered to patients in an invasive fashion via injection of an aqueous drug solution up to four times daily, generally by subcutaneous route or by a long acting depot polymer based formulation that is injected every 3-4 weeks with supplemental injections of aqueous drug as needed.

Thus, treatments using currently available somatostatin peptide-based therapeutics lack ease of administration and are not highly selective to specific somatostatin receptor subtypes. Therefore, there exists an unfulfilled need for longer acting, therapeutic non-peptide somatostatin receptor ligands that have greater receptor subtype selectivity, can readily penetrate the BBB and/or the BRB, and can be delivered using non-invasive pharmaceutical methods including oral administration. It is the object of this invention to provide non-peptide somatostatin receptor ligands that meet the unfulfilled needs detailed above.

Brief Summary of the Invention

The subject invention provides non-peptide compounds which are capable of binding to somatostatin receptors. In one embodiment of this invention, these compounds bind to, and activate, a somatostatin receptor. Particularly preferred compounds are those which selectively bind to and activate SSTR2 receptors. The preferred compounds of the subject invention are particularly advantageous because these compounds selectively bind to somatostatin receptors, are stable, are readily administered, and in a specific embodiment, are able to penetrate the BBB and the BRB.

One embodiment of the subject invention provides potent, non-peptide somatostatin receptor ligands based on a central imidazolin-2-one, imidazolin-4-one, or imidazoline-2,4-dione ring as a core heterocyclic scaffolding structure. Appended to the heterocyclic scaffold are three side chains which account for the advantageous activity of the compounds. In a preferred embodiment, at least two groups appended to the core heterocyclic structure contain one or more specific structural elements or substructures that restrict the conformational mobility of these appended groups. In one embodiment, side chains fix this ligand structure in a 3-dimensional geometry that is useful in

promoting beneficial biological activity. For example, such ligand structures provided herein elicit biological effects similar to somatostatin upon binding to somatostatin receptors.

5 Compounds of the present invention may contain one or more asymmetric carbon atoms and may exist in a given diastereomeric, racemic, or enantiomeric form. In the case of the asymmetric carbon atom C5, as labeled in Formulas IA, IB, and IC, it is generally preferred for the somatostatin agonist to have the *S*-stereochemical assignment. This configuration corresponds to the absolute stereochemistry as derived from *L*-amino acid derivatives (*e.g.*, *L*-(+)-tryptophan) as starting materials from which the instant
10 compounds may be prepared. In one embodiment, compounds of the present invention have the *S*-stereochemical assignment at C5 of Formulas IA, IB, and IC.

In one embodiment of the subject invention, the compounds act as agonists highly specific for the SSTR2 receptor subtype. Also, specifically exemplified herein are somatostatin receptor ligands which are small and exhibit superior penetration of the
15 BBB and BRB. Consequently, the somatostatin receptor ligands of the subject invention are useful for a broad range of therapeutic purposes. These include disorders of the endocrine, gastrointestinal, CNS, PNS, vascular and immune systems as well as cancer.

One category of therapeutic use for the compounds of the subject invention is the treatment of tumors and other forms of cancer. In one embodiment, somatostatin receptor
20 ligands of the present invention can be used to treat solid tumors by efficiently inhibiting angiogenesis. In addition, compounds of the invention retard tumors bearing somatostatin receptors by inhibiting tumor cell growth and proliferation. The ability of compounds of the subject invention to treat various kinds of tumors can be readily confirmed using standard tests including, but not limited to, proliferation tests with
25 various cancer cell lines and tumor growth experiments in nude mice with hormone dependent tumors (see Weckbecker *et al.*, *Cancer Research* 1994, 54: 6334-6337). Thus, the compounds can be used in the treatment of, for example, cancers of the breast, the prostate, the colon, the pancreas, the brain, and the lung.

Another object of the present invention is to provide unique therapeutic
30 compounds that aid in inhibiting gastric, exocrine, and endocrine pancreatic secretion and the release of various peptides of the gastrointestinal tract. These compounds can also be used in the treatment of endocrinological disorders associated with an excess of hormone release, such as growth hormone, glucagon, and insulin.

It is another object of the present invention to provide compounds formulated for oral administration. For example, solutions or suspensions of these compounds may be formulated in the form of tablets, capsules (including time release and sustained release formulations), or pills.

5 Thus, in one embodiment, the present invention provides compounds for use in the treatment of gastro-intestinal (GI) disorders, for example in the treatment of peptic ulcers, disturbances of GI motility, enterocutaneous and pancreaticocutaneous fistula, irritable bowel syndrome, intestinal obstruction, dumping syndrome, watery diarrhea syndrome, acute pancreatitis and gastro-intestinal hormone secreting tumors (*e.g.*,
10 vipomas, glucagonomas, insulinomas, carcinoids and the like).

 The present invention also provides novel somatostatin receptor ligands for use in the treatment of multiple sclerosis, pain, anxiety, depression, schizophrenia, neurodegenerative diseases such as Alzheimer's disease and dementia, and epilepsy. In further embodiments, the compounds of the subject invention can be used in the
15 treatment of vascular disorders. Vascular disorders which can be treated include variceal bleeding, and graft vessel diseases such as restenosis or vascular occlusion following vascular insult (*e.g.*, angioplasty, allo- or xenotransplant vasculopathies; graft vessel atherosclerosis, and transplantation of an organ).

 Another object of the present invention is to provide compounds useful in the
20 treatment of diabetes and diabetes-related pathologies, including angiopathy, dawn phenomenon, neuropathy, nephropathy, and retinopathy (Grant *et al.*, *Diabetes Care* 2000, 23, 504-509).

 Another object of the present invention is to provide the use of novel somatostatin receptor ligand compounds for the manufacture of a medicament for the treatment of
25 conditions as set forth herein.

 In still a further object, the invention provides methods for the treatment of the conditions mentioned above, in a patient in need of such treatment, which comprises administering to such patient a therapeutically effective amount of a somatostatin receptor ligand compound of the subject invention.

30 The present invention provides compounds that may be administered in free form or in pharmaceutically acceptable salt form or complexes. Such salts and complexes may be prepared in a conventional manner and exhibit the activity of the free compounds. A related object of the present invention is to provide pharmaceutical compositions

comprising somatostatin receptor ligands in free base form or in pharmaceutically acceptable acid addition salt form in association with a pharmaceutically acceptable diluent or carrier.

5

Brief Description of the Drawings

Figure 1 illustrates the binding affinity of Compound Example 15a to human SSTR2 receptors according to the present invention.

Figure 2 illustrates the binding affinity of Compound Example 15b to human SSTR2 receptors according to the present invention.

10

Detailed Description of the Invention

The subject invention provides non-peptide compounds that are capable of binding to somatostatin receptors. In one embodiment of this invention, these compounds bind to, and activate, a somatostatin receptor. Preferred compounds of the subject invention are those compounds which selectively bind to somatostatin receptors, are stable, are readily administered, and in a specific embodiment, are able to penetrate the BBB and the BRB. Particularly preferred compounds are those which selectively bind to and activate somatostatin subtype 2 (SSTR2) receptors. The compounds are useful in the treatment of acromegaly, diarrhea management, diabetic retinopathy, macular degeneration, osteosarcoma, irritable bowel syndrome, and for the treatment of abnormal cellular proliferation, including tumors and cancer, and in particular, neuroendocrine tumors and breast cancer.

15

20

In another embodiment, the compounds may treat certain of these disorders through a mechanism that does not involve a somatostatin receptor.


25

One embodiment of the subject invention provides potent, non-peptide somatostatin receptor ligands based on a central imidazolin-2-one, imidazolin-4-one or imidazoline-2,4-dione ring as a core heterocyclic scaffolding structure. Appended to the heterocyclic scaffold are at least three side chains that account for the advantageous activity of the compounds. In a preferred embodiment, at least two groups appended to the core heterocyclic structure include one or more structural elements that restrict the conformational mobility of these appended groups. This ligand structure fixes side chains in the 3-dimensional geometry necessary for biological activity when bound by a

30

somatostatin receptor. Such ligand structures provided herein elicit biological effects similar to somatostatin upon binding to somatostatin receptors.

In certain embodiments of the subject invention, novel somatostatin receptor ligands of Formula IA are provided as follows:




Formula IA

or a pharmaceutically acceptable salt or prodrug thereof, wherein


A and B independently are O or (H, H);

R₁ is a group of formula



wherein the R_a groups are, independently, hydrogen, alkyl, hydroxyalkyl, alkoxyalkyl, or alkylaryl; X is aryloxy, cyano, -CO₂R_b, or -C(O)N(R_c)₂ wherein R_b is H, alkyl or alkylaryl and wherein the R_c groups are, independently, hydrogen, alkyl, hydroxyalkyl, alkoxyalkyl, aminoalkyl, alkylaryl, heterocyloalkyl or are optionally joined to form a six membered ring, including for example, morpholine, piperidine, or piperazine ring, optionally substituted with an alkyl group; Q is an alkyl spacer group optionally interrupted by oxygen, sulfur or nitrogen atoms and optionally substituted by hydroxy, alkoxy, alkyl, or aryl; Z₁, Z₂, and Z₃ are, independently, carbon, oxygen, sulfur, or nitrogen optionally substituted with an alkyl; n is an integer from 0 to 2; and Y is either hydrogen, an alkyl, heterocyloalkyl, or CH₂N(R_a)₂;


R_2 is $-\text{CH}(R_d)-\text{Ar}$ or $-\text{SO}_2-\text{Ar}$ wherein R_d is alkyl, aryl, cyano, $-\text{CO}_2R_b$, or $-\text{C}(\text{O})\text{N}(R_c)_2$; Ar is phenyl or naphthyl optionally and independently, mono- or di-substituted by hydroxy, cyano, halogen, alkyl, alkoxy, $-\text{N}(R_a)_2$, alkylthio, alkylsulfonyl, arylsulfonyl, nitro, $-\text{S}(\text{O})_2\text{N}(R_c)_2$, trifluoromethyl,



$-\text{CH}_2\text{N}(R_a)_2$, $-\text{CO}_2R_b$, $-\text{C}(\text{O})\text{N}(R_c)_2$, or an aromatic group of formula wherein A is CH_2 , O, NH, S or CO and the R_e groups, independently, are hydrogen, alkyl, alkoxy, trifluoromethyl, cyano, nitro, amino, halogen, hydroxy;

R_3 is selected from hydrogen, alkyl, aryl, alkylaryl or heteroaryl; and

R_4 is a group of formula:



wherein Z_1 , Z_2 , and Z_3 are, independently, carbon, oxygen, sulfur or nitrogen optionally substituted with an alkyl; R_f is hydrogen, alkyl or an aryl group; and R_g is hydrogen, halogen, cyano, trifluoromethyl, aryl, alkyl or alkoxy.

In a preferred embodiment of Formula IA, A and B are oxygen.

R_1 of Formula IA preferably is a group in which 5 or 6 atoms, at least four of which are carbon, separate a basic nitrogen atom from the point of attachment.


R_2 of Formula IA preferably is a group of formula $-\text{SO}_2-\text{Ar}$ wherein Ar is an optionally substituted diphenyl ether structure.

R_3 of Formula IA preferably is H.

R_4 of Formula IA preferably is an optionally substituted 3-indolyl group.

Absolute stereochemistry at C5 of Formula IA preferably is S.

In certain other embodiments of the subject invention, novel somatostatin receptor ligands of Formula IB are provided as follows:




Formula IB

or a pharmaceutically acceptable salt or prodrug thereof, wherein

5 A and B independently are O or (H, H);

Z₁, Z₂, and Z₃ are independently carbon, oxygen, sulfur or a nitrogen atom optionally substituted with an alkyl group;


R₁ is a group of formula



wherein X is a saturated or unsaturated aliphatic C₂₋₄ carbonic chain which is (a) optionally interrupted by -O- or -S- or -N(R_b)- and/or (b) optionally substituted by hydroxy, alkoxy groups, alkyl, or aryl; and wherein R_a are independently hydrogen, alkyl, or together form a 3-6

carbon-membered ring; and the R_b independently are hydrogen, alkyl, or arylalkyl; and n is an integer from 0 to 2;

R_2 is a group of formula $-SO_2-Ar$ or $-C(R_c)H-Ar$ wherein R_c is alkyl, or phenyl optionally substituted by up to three groups alternatively from hydroxy, halogen, alkyl, alkoxy, cyano, trifluoromethyl, amino, monoalkylamino, dialkylamino, carboxy, carboxyalkyl, monoalkylamino-carbonyl, dialkylaminocarbonyl, alkylthio; Ar is phenyl or naphthyl optionally mono- or di-substituted by hydroxy, halogen, alkyl, alkoxy, cyano, trifluoromethyl, aminomethyl, monoalkylaminocarbonyl, dialkylaminocarbonyl, morpholinocarbonyl, piperazinocarbonyl, or a group of formula



wherein A is CH_2 , O , S or CO ; R_d independently are hydrogen, alkyl, alkoxy, trifluoromethyl, amino, halogen, or hydroxy;

R_3 is hydrogen, alkyl, aryl, heteroaryl or alkylaryl;

R_4 is alkyl, aryl, or heteroaryl; and

R_5 are independently hydrogen, halogen, cyano, alkyl or alkoxy.

In a preferred embodiment of Formula IB, A and B are O .

Z_1 and Z_2 of Formula IB are preferably carbon atoms connected by a double bond and Z_3 is preferably $N-H$ (e.g., indole).

R_1 of Formula IB preferably is a group in which 5 atoms separate a basic nitrogen atom from the point of attachment.


R_2 of Formula IB preferably is a group of formula $-SO_2-Ar$ wherein Ar is an optionally substituted diphenyl ether structure.

R_3 of Formula IB preferably is hydrogen or methyl.

R_4 of Formula IB is preferably methyl or substituted phenyl.

R_5 of Formula IB is preferably hydrogen, methyl or halogen.

In a preferred embodiment, the subject invention provides novel somatostatin receptor ligands of Formula IC




Formula IC

or a pharmaceutically acceptable salt or prodrug thereof, wherein:


5 A and B independently are oxygen, or (H,H);

R_1 is a group selected from the following Formula Set A or Formula Set B



Formula Set A

10



Formula Set B

wherein;

15 each R_a independently is hydrogen, alkyl, alkenyl, alkynyl, hydroxyalkyl, alkoxyalkyl, arylalkyl, or form together a 3-6 carbon-membered ring;

G is aryloxy, cyano, $-\text{CO}_2R_a$, or $-\text{C}(\text{O})\text{N}(\text{R}_b)_2$ wherein R_b is independently hydrogen, alkyl, alkenyl, alkynyl, hydroxyalkyl, alkoxyalkyl, monoalkylaminoalkyl, dialkylaminoalkyl, aminoalkyl, arylalkyl, heterocyclalkyl, or is optionally joined to a five- or six-membered ring,

including for example pyrrolidine, morpholine, piperidine, or piperazine ring, optionally substituted with an alkyl group;

Q is a C₁-C₅ alkyl, alkenyl, or alkynyl linker group that can optionally include oxygen, sulfur, or nitrogen atoms or -N(R_a)- and/or is optionally substituted by hydroxy, alkoxy, alkyl, or aryl;

each E₁, E₂, and E₃ independently is a carbon, oxygen, sulfur, covalent bond, or nitrogen optionally substituted with an alkyl, alkenyl, alkynyl, alkylaryl, or aryl;

n is an independently 0, 1, or 2; and


J is hydrogen, alkyl, heterocyclalkyl, or CH₂N(R_a)₂;

R₂ is -SO₂-Ar or (CHR_c)_n-Ar

wherein

R_c is hydrogen, alkyl, alkenyl, alkynyl, hydroxy, alkoxy, hydroxyalkyl, alkoxyalkyl, aryl, aryloxy, cyano, -CO₂R_a or -C(O)N(R_b)₂;

Ar is phenyl or naphthyl, optionally and independently mono- or disubstituted with hydroxy, cyano, halogen, alkyl, alkenyl, alkynyl, alkoxy, alkylthio, alkylsulfonyl, arylsulfonyl, amino, nitro, trifluoromethyl, aminomethyl, dimethylaminocarbonyl, benzimidazolyloxy, morpholinocarbonyl, or an aromatic group of formula:



wherein D is oxygen, sulfur, CH₂, or CO; and each R_d is independently hydrogen, alkyl, alkenyl, alkynyl, alkoxy, trifluoromethyl, amino, halogen, or hydroxy;


R₃ is selected from hydrogen, alkyl, alkenyl, alkynyl, aryl, or heteroaryl; and

R₄ is either a group of formula -CH₂-Y or CHR_e-Y if R₁ is a group of Formula Set A or a group of formula CHR_e-Y if R₁ is a group of Formula Set B;

wherein

R_e is alkyl, alkenyl, alkynyl, hydroxyalkyl, alkoxyalkyl, aminoalkyl, heterocyclalkyl;

Y is a group of formula



wherein R_f is hydrogen, halogen, or alkoxy; and R_g is hydrogen, alkyl, or benzyl.


5 In a preferred embodiment of Formula IC, A and B are oxygen.

In one embodiment of Formula IC, A and B are oxygen; R_1 preferably is a group of Formula Set A as defined above wherein four to six atoms separate a basic nitrogen atom from the point of attachment at N3 of the heterocyclic core structure of Formula IC; R_2 preferably is a group of formula SO_2 -Ar wherein Ar is an optionally substituted diphenyl ether structure; R_3 preferably is hydrogen; and R_4 is as defined above wherein R_e preferably is hydrogen, methyl, or substituted phenyl, Y is an optionally substituted indole ring, R_f preferably is hydrogen, methyl, or halogen, and R_g preferably is hydrogen.

15 In an alternate preferred embodiment of Formula IC, A and B are oxygen; R_1 preferably is a group of Formula Set B as defined above wherein four to six atoms separate a basic nitrogen atom from the point of attachment at N3 of the heterocyclic core structure of Formula IC; R_2 preferably is a group of formula SO_2 -Ar wherein Ar is an optionally substituted diphenyl ether structure; R_3 preferably is hydrogen or methyl; and R_4 is as defined above wherein R_e preferably is methyl or substituted phenyl, Y preferably is an optionally substituted indole ring, R_f preferably is hydrogen, methyl, or halogen, and R_g preferably is hydrogen.

20 In a further embodiment of Formula IC, various structural elements of the R_1 - R_4 groups restrict the conformational mobility of two or more of the R_1 - R_4 groups appended to the heterocyclic core structure of Formula IC. By restricting the conformation mobility of certain R_1 - R_4 groups, the structure of Formula IC is biased in a desired 3-dimensional geometry which promotes beneficial biological activity and/or potency/selectivity for a given somatostatin receptor subtype. In a preferred embodiment, the structural elements are, independently any of (a), (b) or (c): (a) a group of formula $-SO_2-$ (sulfonyl group) attached directly either to the N1 or N3 of the heterocyclic core structure of Formula IC;

or (b) a branched group having the structure



wherein each R_h independently is any suitable carbon substituent consistent with the structure for Formula IC (*e.g.*, C_1 - C_{10} alkyl), and such branched group is attached directly to either N1 or C5 of the heterocyclic core structure of Formula IC; or (c) a 1,3-cis-disubstituted cyclohexyl group embedded within the R_1 group appended to N3 of the heterocyclic core structure of Formula IC.

Compounds of the present invention may contain one or more asymmetric carbon atoms and may exist in a given diastereomeric, racemic, or enantiomeric form. In a preferred embodiment, the active compound has an absolute *S*-stereochemical attachment at C5 of Formulas IA, IB, and IC.

Definitions for Structures of Formula IA and Formula IB

The following definitions are applicable to those structures as defined above for Formula IA and Formula IB.

Unless otherwise specified, as used herein, the term "alkyl" refers to a straight or branched chain alkyl moiety. In one embodiment, the alkyl moiety is C_{1-8} alkyl, which refers to an alkyl moiety having from one to eight carbon atoms, including for example, methyl, ethyl, propyl, isopropyl, butyl, *tert*-butyl, pentyl, hexyl, and octyl.

The term "alkyl" also includes cycloalkyl, including for example cyclohexyl and the like. The term "alkyl" also includes alkenyl, which refers to a straight or branched alkyl moiety having one or more carbon double bonds, of either *E* or *Z* stereochemistry where applicable, and includes for example, vinyl, 1-propenyl, 1- and 2- butenyl, and 2- methyl-2-propenyl.

The term "alkyl" also refers to an alicyclic moiety having from three to six carbon atoms and having in addition one double bond. Such groups, also known as "cycloalkenyl," include, for example, cyclopentenyl and cyclohexenyl.

The term "alkyl" also includes alkynyl moieties, including for example, ethynyl, 1-propynyl, 1- and 2- butynyl, 1- methyl-2-butyne, and the like.

The term "alkoxy" refers to an alkyl-O-group, in which the alkyl group is as described within this section for the structures of Formula IA and Formula IB.

The term "halogen" is refers to fluorine, chlorine, bromine and iodine.

The term "aryl" refers to an aromatic carbocyclic ring, *e.g.*, phenyl, substituted phenyl and like groups, as well as rings which are fused, *e.g.*, naphthyl, biphenyl indaryl and the like. Aryl groups may be optionally substituted with from 1 to 3 groups of alkyl, aryl, heteroaryl, halogen, hydroxy, alkoxy, nitro, cyano, trifluoromethyl, alkylthio, alkylsulfonyl, arylsulfonyl, $N(R_a)_2$, $S(O)_2N(R_c)_2$, $CH_2N(R_a)_2$, CO_2R_b , and $C(O)N(R_c)_2$.

The term "arylalkyl" refers to a moiety in which the "aryl" and "alkyl" groups are as described within this section for the structures of Formula IA and Formula IB.

The term "heterocycloalkyl" refers to a saturated heterocyclic moiety having from two to six carbon atoms and one or more heteroatom from the group N, O, and S (or oxidized versions thereof) which may be optionally benzofused at any available position. This includes, for example, azetidiny, pyrrolidiny, tetrahydrofuranyl, piperidiny, benzodioxolyl and the like.

The term "heterocycloalkyl" also refers to an alicyclic moiety having from three to six carbon atoms and one or more heteroatoms from the group N, O, and S and having in addition one double bond. Such moieties may also be referred to as "heterocycloalkenyl" and includes, for example, dihydropyranyl, and the like.

In the case that the heterocyclyl moiety contains a nitrogen atom that may be substituted, then a substituent may be chosen from the group of R_b .

The term "heteroaryl" refers to monocyclic or bicyclic aromatic ring systems of five to ten atoms of which at least one atom is selected from O, N, and S, in which a carbon or nitrogen atom is the point of attachment, and in which one additional carbon atom is optionally replaced with a heteroatom selected from O or S, and in which from 1 to 3 additional carbon atoms are replaced by nitrogen heteroatoms. The heteroaryl group is optionally substituted with up to three groups selected from alkyl, aryl, heteroaryl, halogen, hydroxy, alkoxy, nitro, cyano, trifluoromethyl, alkylthio, alkylsulfonyl, arylsulfonyl, $N(R_a)_2$, $S(O)_2N(R_c)_2$, $CH_2N(R_a)_2$, CO_2R_b , and $C(O)N(R_c)_2$. This term includes, for example, imidazolyl, tetrazolyl, furanyl, thiophenyl, pyridyl, indolyl, quinolyl, benzothiadiazolyl, benzofurazanyl, benzotriazolyl, and the like.

Certain of the above defined terms may occur more than once in Formula IA and/or Formula IB, and upon such occurrence each term shall be defined independently of the other.

The term "orthogonal protecting group" refers to a protecting group which can be removed from a compound while leaving other protecting groups in the compound intact.

Definitions for Structures of Formula IC

The following definitions are applicable to those structures as defined above for Formula IC.

5 Unless otherwise specified, as used herein, the term "alkyl" refers to a straight, branched chain, or cyclic alkyl moiety containing 1-15 carbon atoms. Preferred straight or branched alkyl groups include methyl, ethyl, propyl, isopropyl, butyl, *tert*-butyl, pentyl, hexyl, octyl. Preferred cycloalkyl groups include, cyclopentyl and cyclohexyl. The alkyl group specifically includes fluorinated alkyls such as CF₃ and other halogenated alkyls
10 such as CH₂CF₂, CF₂CF₃, the chloro analogs, and the like.

Unless otherwise specified, the term "acyl," as used herein, refers to a group of the formula C(O)R', wherein R' is an alkyl, aryl, arylalkyl group, wherein these groups are as defined within this section for the structures of Formula IC.

The term "alkenyl" refers to a straight or branched alkyl moiety having one or
15 more carbon double bonds, of either *E* or *Z* stereochemistry where applicable. This term includes for example, vinyl, 1-propenyl, 1- and 2- butenyl, and 2- methyl-2-propenyl, as well as "cycloalkenyl" groups such as cyclopentenyl and cyclohexenyl.

The term "alkynyl" refers to a straight, branched or cyclic alkyl moiety with at least one triple bond, including but not limited to ethynyl, 1-propynyl, 1- and 2- butynyl,
20 1- methyl-2-butynyl groups and the like.

The term "alkoxy," as used herein, and unless otherwise specified, refers to a moiety of the structure -O-alkyl, -O-alkenyl, or -O-alkynyl wherein alkyl, alkenyl and alkynyl are as defined above. The alkyl group can be optionally substituted as described above. Alkoxy groups can include methoxy, ethoxy, propoxy, isopropoxy, butoxy,
25 isobutoxy, tertiary butoxy, allyloxy, propargyloxy, OCF₃, OCH₂CF₃, OCF₂CF₃ and the like.

The term "alkylthio," as used herein, refers to an alkyl group attached to the molecule through a sulfur atom.

The term "aryl," as used herein, and unless otherwise specified, refers to either
30 aromatic carbocyclic ring or rings which are fused including, for example, phenyl, biphenyl, or naphthyl. The aryl group can be optionally substituted with one or more of the moieties selected from the group consisting of alkyl, heteroaryl, heterocyclic, alkoxy, aryloxy, arylalkoxy, heteroarylalkoxy, carbohydrate, amino acid, amino acid esters,

amino acid amides, halo, hydroxyl, carboxyl, acyl, acyloxy, nitro, cyano, thiol, imide, sulfonic acid, sulfate, sulfonyl, sulfanyl, sulfinyl, sulfamoyl, carboxylic ester, carboxylic acid, amide, phosphonyl, phosphinyl, phosphoryl, thioester, thioether, oxime, hydrazine, carbamate, phosphonic acid, phosphate, phosphonate, phosphinate, hydroxamic acid, sulfonylimide, $-N(R_a)_2$, $-N(R_a)C(O)R_a$, $-N(R_a)C(O)N(R_a)$, $-N(R_a)S(O)_2R_a$, $-S(O)_2N(R_a)_2$, $-CH_2N(R_a)_2$, $-CO_2R_a$, and $-C(O)N(R_a)_2$, optionally substituted with, or fused with, an aryl group. This term includes, for example phenyl or naphthyl or any other desired functional group that does not inhibit the pharmacological activity of this compound. Alternatively, adjacent groups on the aryl ring may combine to form a five- to seven-membered carbocyclic, aryl, heteroaryl or heterocyclic ring. In another embodiment, the aryl ring is substituted with an optionally substituted cycloalkyl (such as cyclopentyl or cyclohexyl), or an alkylene dioxy moiety (for example methylenedioxy).

The term "arylalkyl," as used herein, and unless otherwise specified, refers to an aryl group as defined above linked to the molecule through an alkyl group as defined above. The aryl portions can be optionally substituted as described within this section for the structures of Formula IC.

The term "aryloxy," as used herein, refers to an aryl or heteroaryl group bound to the molecule through an oxygen atom. The aryl group can be optionally substituted as set out within this section (for the structures of Formula IC) for aryl groups. Preferred aryloxy groups include optionally substituted phenoxy and pyridinyloxy.

The term "halogen" refers to halogen atoms including, for example, fluorine, chlorine, bromine, and iodine.

The terms "heteroaryl" and "heteroaromatic," as used herein, refer to monocyclic or bicyclic aromatic ring systems of five to ten atoms of which at least one atom is selected from O, N, and S, in which a carbon or nitrogen atom is the point of attachment, and in which one additional carbon atom is optionally replaced with a heteroatom selected from O or S, and in which from 1 to 3 additional carbon atoms are replaced by nitrogen heteroatoms. In one embodiment, the heteroaryl group is optionally substituted with up to three groups selected from 1 to 3 of from alkyl, aryl, heteroaryl, heterocyclyl, halogen, hydroxy, alkoxy, nitro, cyano, trifluoromethyl, alkylthio, alkylsulfonyl, arylsulfonyl, $-N(R_a)_2$, $-N(R_a)C(O)R_a$, $-N(R_a)C(O)N(R_a)$, $-N(R_a)S(O)_2R_a$, $-S(O)_2N(R_a)_2$, $-CH_2N(R_a)_2$, $-CO_2R_a$, and $-C(O)N(R_a)_2$, optionally substituted with, or fused with, an aryl group. This term (heteroaryl) includes, for example, phenyl or naphthyl.

Heteroaryl thus includes aromatic and partially aromatic groups which contain one or more heteroatoms. Examples of this type include but are not limited to are furan, benzofuran, thiophene, benzothiophene, pyrrole, pyrazole, imidazole, oxazole, benzoxazole, thiazole, benzthiazole, isothiazole, thiadiazole, triazole, benzotriazole, furazan, benzofurazan, thiafurazan, benzothiafurazan, tetrazole, oxadiazole, triazine, pyridine, pyridazine, pyrimidine, pyrazine, triazine, indolizine, indole, isoindole, purine, quinoline, benzimidazole, pteridine, isoquinoline, cinnoline, quinazoline, quinoxaline,

The term "heteroarylalkyl," as used within this section for the structures of Formula IC, refers to a heteroaryl group, as defined above in this section, linked to the molecule through an alkyl group as defined above within this section.

The term "heterocyclyl" and "heterocyclic," as used herein, refers to a nonaromatic cycloalkyl group that may be partially (contains at least one double bond) or fully saturated and wherein one of the carbon atoms in the ring is replaced by a heteroatom, selected from O, S, SO, SO₂, or N, and in which up to three additional carbon atoms may be optionally replaced with heteroatoms. Heterocyclic rings are monocyclic, or are fused, bridged, or spiro polycyclic ring systems. Monocyclic heterocyclic rings contain from 3 to 9 member atoms (carbon and heteroatoms), and preferably from 5 to 7 member atoms, in the ring. Polycyclic heterocyclic rings contain from 7 to 12 member atoms in the ring. Preferred polycyclic heterocyclic rings comprise 5-, 6- or 7-membered rings fused to 5-, 6- or 7-membered rings.

Heterocyclyl is linked to a carbon or nitrogen atom. If the heterocyclyl is carbon linked and contains an additional nitrogen, then the additional nitrogen may be substituted by R_a. If the heterocyclyl is nitrogen linked and contains an additional nitrogen, then the additional nitrogen may be substituted by R_a. Examples of heterocyclyls are piperidinyl, morpholinyl, pyrrolidinyl, tetrahydrofuranyl, dihydrofuranyl, tetrahydropyranyl, dihydropyranyl, imidazolyl, piperazinyl, pyrrolidin-2-onyl, piperidin-2-onyl, succinimidonyl, imidazolidinonyl and the like.

The term "heterocyclylalkyl," as used herein, refers to a heterocyclyl group as defined above linked to the molecule through an alkyl group as defined above.

The term "protected" or "protecting group" as used herein and unless otherwise defined refers to a group that is added to an oxygen, nitrogen, sulfur or phosphorus atom

to prevent its further reaction or for other purposes. A wide variety of oxygen and nitrogen protecting groups are known to those skilled in the art of organic synthesis.

General Definitions

5 The term "mammal" includes humans and all domestic and wild animals including, without limitation, cattle, horses, swine, sheep, goats, dogs, cats, rabbits and the like.

 "Optional" or "optionally" means that the subsequently described event, element, or circumstance may or may not occur, and that the description includes instances where
10 said event occurs and instances in which it does not.

 The term "enantiomer," as used herein unless otherwise specified, refers to a compound of the present invention having one or more chiral carbon atoms with absolute *S* or *R* stereochemical configuration. The term "counter enantiomer" refers to the isomer in which all carbon atoms have the opposite absolute stereochemical configuration. The
15 term "enantiomerically enriched," as used herein, refers to an active compound substantially in the absence of its counter enantiomer. The term "enantiomerically pure," as used herein, refers to a composition having active compounds in excess of 98% absolute stereochemical purity.

 Salts encompassed within the terms "pharmaceutically acceptable salts" or
20 "acceptable acid addition salts" refer to non-toxic salts of the compounds of this invention which are generally prepared by reacting the free base with a suitable organic or inorganic acid. Representative salts include, but are not limited to, the following: acetate, benzenesulfonate, benzoate, bicarbonate, bisulfate, bitartrate, borate, camsylate, carbonate, citrate, dihydrochloride, edetate, edisylate, estolate, esylate, fumarate, gluconate, glutamate, hydrobromide, hydrochloride, hydroxynaphthoate, lactate,
25 lactobionate, laurate, malate, maleate, mandelate, mesylate, mucate, napsylate, nitrate, N-methylglucamine ammonium salt, oleate, oxalate, palmoate (embonate), palmitate, pantothenate, phosphate/diphosphate, polygalacturonate, salicylate, stearate, sulfate, subacetate, succinate, tannate, tartrate, tosylate, and valerate.

30 The term "substituted" shall be deemed to include multiple degrees of substitution by a named substituent. Where multiple substituent moieties are disclosed or claimed, the substituted compound can be independently substituted by one or more of the disclosed or claimed substituent moieties, singly or severally.

As used herein, the term “substantially free of” or “substantially in the absence of” or “isolated” refers to a composition that includes at least 95% to 98% by weight, and even more preferably 99% to 100% by weight, of the designated enantiomer of that compound of the present invention. The compounds of the present invention typically have at least one chiral carbon, and therefore, exist as enantiomers. In a preferred embodiment, the active compound is used substantially in the absence of its counter enantiomer.

The term “treatment” as used herein covers any treatment of a disease or disorder in a mammal, particularly a human and includes:

- I. Preventing the disease from occurring in a subject which may be predisposed to the disease but has not yet been diagnosed as having it;
- II. Inhibiting the disease, *e.g.*, arresting its development; or
- III. Relieving the disease, *e.g.*, causing regression of the disease.

Biological Activity and Therapeutic Applications

The somatostatin receptor ligand compounds of the subject invention and their analogs, derivatives, and physiologically acceptable salts, have highly advantageous pharmacological properties. Of particular importance is the ability of these compounds to bind to human somatostatin receptors. In a preferred embodiment, compounds of the subject invention are selective agonists for somatostatin SSTR2 receptors. This activity can be confirmed by, for example, somatostatin receptor binding studies and second messenger functional assays for somatostatin receptors (see Kaupmann *et al.*, *FEBS Letters* 1993, 331, 53-59). In a preferred embodiment, the compounds of this invention inhibit the binding of somatostatin to its receptor at an IC_{50} of about 100 pM to about 1 μ m.

The *in vitro* biological activity of the compounds of the subject invention can be confirmed *in vivo* utilizing any of a number of assays well known to those skilled in the art. For example, the compounds can be evaluated for their ability to depress serum GH and insulin levels in the rat. In one example, the test is carried out using male rats. The test substance is administered at varying, logarithmically staggered doses preferably employing at least five (5) rats per dose. One hour after administration of the test substance, blood is taken. The determination of the blood serum GH and insulin levels is measured by radio-immunoassay.

The ability of the compounds of the present invention to act as somatostatin agonists makes them useful as pharmacologic agents for mammals, especially for humans, for the treatment and prevention of disorders wherein somatostatin itself, or the physiological processes it regulates, are involved. These include disorders of the endocrine, gastrointestinal, CNS, PNS, vascular and immune systems as well as cancer.

The somatostatin receptor ligands of the present invention can be used to inhibit various hormone secretions and trophic factors in mammals. Thus, the somatostatin receptor ligands of the subject invention may be administered to treat disorders involving, for example, autocrine or paracrine secretions of trophic factors including cancers of the breast, brain, prostate, and lung (both small cell and non-small cell epidermoids), as well as hepatomas, neuroblastomas, colon and pancreatic adenocarcinomas (ductal type), chondrosarcomas, and melanomas. The compounds of the subject invention can be used to treat SSSTR-2 receptor bearing tumors. This activity can be confirmed in proliferation tests with various different cancer cell lines and in tumor growth experiments in nude mice with hormone dependent tumors (see Weckbecker *et al.*, *Cancer Research* 1994, 54: 6334-6337).

In one embodiment, somatostatin receptor ligands of the present invention promote apoptosis and can be used to treat cancer directly or to sensitize cancer cells for combination treatments using other regimens including radiation therapy or chemotherapy.

Uses of these compounds also include suppressing certain endocrine secretions, such as, insulin, glucagons, prolactin, and GH, which in turn can further suppress the secretion of various trophic factors such as IGF-1. The compounds of the invention are accordingly indicated for use in the treatment of disorders with an etiology comprising or associated with excess GH and trophic factor secretion. The ability to suppress these secretions is useful in the treatment of disorders such as acromegaly. This activity is also useful in the treatment of neuroendocrine tumors, such as carcinoids, vipomas, insulinomas and glucagonomas. The compounds of this invention are also useful for treating diabetes and diabetes-related pathologies, including angiopathy, dawn phenomenon, neuropathy, nephropathy, and retinopathy (Grant *et al.*, *Diabetes Care* 2000, 23, 504-509).

In another embodiment, compounds of the subject invention can be used to treat vascular disorders including bleeding disorders of the gastrointestinal system, such as

those involving the splanchnic blood flow and esophageal varices associated with diseases such as cirrhosis. The ability of the compounds of to mediate vasoconstriction also render them useful in the treatment of cluster headache and migraine.

5 Compounds of the invention can also be used to inhibit the proliferation of vascular endothelial cells and so are indicated for use in treating graft vessel diseases such as restenosis or vascular occlusion following vascular insult such as angioplasty, allo- or xenotransplant vasculopathies, graft vessel atherosclerosis, and in the transplantation of an organ (*e.g.*, heart, liver, lung, kidney or pancreatic transplants (Weckbecker *et al.*, *Transplantation Proceedings* 1997, 29, 2599-2600)). Compounds of
10 the invention can also be used to inhibit angiogenesis and are indicated for use in wound healing and treating metastatic stage cancer including but not limited to lung, breast and prostate cancers.

Compounds of the subject invention can also be used for inhibiting gastric and exocrine and endocrine pancreatic secretion and the release of various peptides of the
15 gastrointestinal tract. Thus, these compounds are useful in treating gastro-intestinal disorders, for example in the treatment of peptic ulcers, NSAID-induced ulcers, ulcerative cholangitis, acute pancreatitis (*e.g.*, in post-ERCP patients), enterocutaneous and pancreaticocutaneous fistula, disturbances of GI motility, intestinal obstruction, chronic atrophic gastritis, non-ulcer dyspepsia, scleroderma, irritable bowel syndrome, Crohn's
20 disease, dumping syndrome, watery diarrhea syndrome, and diarrhea associated such diseases as AIDS or cholera (see O'Dorisio *et al.*, *Advances in Endocrinology Metabolism* 1990, 1: 175-230).

Somatostatin receptor ligands of the instant invention can also be used in modulating pro-inflammatory mediators such as substance P and various cytokines
25 (Peluso *et al.*, *Neuropeptides* 1996, 30, 443-451), and may be used in the therapeutic treatment of psoriasis; arthritis; topical inflammation such as is associated with sunburn, eczema, or other sources of allergies, including asthma; and itching.

In a specific embodiment, the compounds disclosed herein are also functional as neuromodulators in the central nervous system, with useful applications in the treatment
30 of neurodegenerative diseases such as stroke, multiple sclerosis, Alzheimer's disease and other forms of dementia, mental health disorders (such as anxiety, depression, and schizophrenia), and in other neurological diseases such as pain and epilepsy (Vezzani *et al.*, *European Journal of Neuroscience* 1999, 11, 3767-3776).

The instant compounds can also be used in combination with other therapeutic agents. For example, in the case of treating organ transplantation, examples of other therapeutic agents include cyclosporin and FK-506. For treating tumors, examples of other agents include tamoxifen and alpha-interferon. For diabetes, examples of other compounds include metformin or other biguanides, acarbose, sulfonylureas thiazolidinediones or other insulin sensitizers including, but not limited to, compounds which function as agonists on peroxisome proliferator-activated receptor gamma (PPAR-gamma), insulin, insulin-like-growth factor I, glucagon-like peptide I (glp-I) and available satiety-promoting agents such as dexfenfluramine or leptin.

Formulations

Compounds of the present invention may be formulated as solutions or suspensions, in the form of tablets, capsules (each including timed release and sustained release formulations), pills, oils, powders, granules, elixers, tinctures, suspensions, syrups, emulsions, microemulsions, or with excipients. Likewise, they may also be administered by any conventional route, for example in intravenous (both bolus and infusion), intraperitoneal, intraocularly, subcutaneous, intramuscular form, enterally, preferably orally (e.g., in the form of tablets or capsules), or in a nasal, buccal, transdermal, or a suppository form, using well known formulations to those of ordinary skill in the pharmaceutical arts.

In addition, the compounds of the present invention can also be administered in the form of liposomes or the like. Disintegrators include, without limitation, delivery systems such as small unilamellar vesicles, large unilamellar vesicles and multilamellar vesicles. Liposomes can be formed from a variety of phospholipids, such as cholesterol, stearylamine or phosphatidylcholines.

For oral administration in the form of a tablet or capsule, the active drug component can be combined with an oral, non-toxic pharmaceutically acceptable inert carrier such as ethanol, glycerol, water and the like. Moreover, when desired or necessary, suitable binders, lubricants, disintegrating agents and coloring agents can also be incorporated into the mixture. Suitable binders include starch, gelatin, natural sugars such as glucose or beta-lactose, corn sweeteners, natural and synthetic gums such as acacia, tragacanth or sodium alginate, carboxymethylcellulose, polyethylene glycol, waxes and the like. Lubricants used in these dosage forms include, for example, sodium

oleate, sodium stearate, magnesium stearate, sodium benzoate, sodium acetate, sodium chloride, starch, methyl cellulose, agar, bentonite, xanthan gum, and the like.

The somatostatin receptor ligand compounds of the subject invention may be administered as prodrugs in which a group including, for example, esters, peptide amides, or Schiff bases of amines, may be metabolically or spontaneously cleaved upon administration and is fixed to the parent biologically active form of the somatostatin receptor ligand.

The dosage regimen for the compounds of the present invention is selected in accordance with a variety of factors including type, species, age, weight, sex and medical condition of the patient; the severity of the condition to be treated; the route of administration; the renal and hepatic function of the patient; and the particular compound or salt thereof employed. An ordinarily skilled physician or veterinarian can readily determine and prescribe the effective amount of the drug required to prevent, counter or arrest the progress of the condition.

In general, satisfactory results in animals are indicated to be obtained at a daily dosage of from about 0.1 to about 200 mg, preferably from about 0.1 to about 5 mg/kg animal body weight. In larger mammals, for example humans, an indicated daily dosage is in the range from about 0.5 to about 100 mg, preferably from about 1 to about 50 mg of an agent of the invention conveniently administered, for example, in divided doses up to four times a day or in sustained release form.

Injected intravenous, subcutaneous or intramuscular dosages of the compounds of the present invention, when used for the indicated effects, will range between about 0.001 to 1.0 mg/kg. Furthermore, preferred compounds for the present invention can be administered in intranasal form via topical use of suitable intranasal vehicles, or via transdermal routes, using those forms of transdermal skin patches well known to those of ordinary skill in that art. To be administered in the form of a transdermal delivery system, the dosage administration can be continuous rather than intermittent throughout the dosage regimen. Transdermal delivery can also be achieved using approaches known to those skilled in the art.

All patents, patent applications, provisional applications and publications referred to or cited herein are incorporated by reference in their entirety to the extent they are not inconsistent with the explicit teachings of this specification.

Synthetic Procedures

Illustrative examples of procedures which can be used to produce the compounds of the subject invention are outlined below.

5 List of Abbreviations:

Ac₂O- acetic anhydride

Alloc- allyloxycarbonyl

Boc- t-butyloxycarbonyl

Cbz- benzyloxycarbonyl

10 CDI- N,N'-carbonyldiimidazole

DCC- dicyclohexylcarbodiimide

DCM- dichloromethane

DIEA-diisopropylethylamine

HOBt- hydroxybenztriazole

15 HPLC-high pressure liquid chromatography

(i-Pr)₂NEt- diisopropylethylamine

LHDMS-lithium bis(trimethylsilylamide)

NMR-nuclear magnetic resonance

Ph-phenyl

20 TBAF-tetrabutyl-ammonium-fluoride

TBDMS-t-butyldimethylsilyl

THF-tetrahydrofuran

THP-tetrahydropyranyl

25 The preparation of compounds of Formula IA, Formula IB, and Formula IC of the present invention may be carried out in sequential or convergent synthetic routes. Compounds with different aromatic or non-aromatic rings and/or bearing additional or different substituents on these rings are readily prepared by minor modification of the methods herein using procedures known in the art.


30 Conditions used to remove protecting groups such as phenolic benzyl ethers may be changed from catalytic hydrogenation in the presence of a noble metal such as palladium supported on charcoal to Lewis acid mediated deprotection, *e.g.*, with BBr₃, in cases where other functional groups are present, *e.g.*, double bonds and aromatic halogen

substituents that are not compatible with catalytic hydrogenation conditions. Synthesis of compounds or intermediates in which multiple amino, hydroxyl and carboxyl group functions require differential protecting group strategies can be carried out by one skilled in the art of organic synthesis by the appropriate selection of orthogonal protecting groups such as benzyl, Cbz, Boc, and Alloc for amino groups and benzyl, methyl, TBDMS, acetyl, THP, for hydroxy groups. Conditions for introducing and removing protecting groups which may be required or present can be found in T. Greene and P. G. M. Wuts, *Protective Groups in Organic Synthesis*, John Wiley & Sons Inc., New York, N. Y. 1991.


Enantiomerically pure or enriched starting materials and intermediates required for the synthesis of enantiomerically pure or enriched compounds of Formulas IA, IB, and IC having defined *R* or *S* stereochemistry at C5 and any other given asymmetric carbon atoms may be prepared from enantiomerically pure or enriched α -amino acids or from enantiomerically pure or enriched hydantoins that may in turn be prepared by established literature methods (Duthaler, *Recent Developments in the Stereoselective Synthesis of α -Amino Acids*, *Tetrahedron* 1994, 50, 1539-1650).

Preparation of Formulas IA, IB, and IC

The compounds of Formula IA, IB, and IC may be prepared as depicted in Scheme I by deprotonating the hydrogen from N1 with a base for a compound of Formula II, wherein A and B are as defined above, wherein the following structure




in Formula IB is hereinafter referred to as R₄, and wherein R'₁, R'₃, and R'₄ are R₁, R₃, and R₄ as defined above or are optionally protected forms of R₁, R₃, and R₄ thereof. By reacting the resulting anion with a compound of Formula III wherein R'₂ is R₂ as defined above or is optionally a protected form of R₂ and V is a reactive leaving group such as trifluoromethanesulfonate, chlorine, bromine, or iodine, the product is deprotected and a compound of Formulas IA, IB, or IC are recovered in free base or acid addition salt form.

Scheme I

Suitable protecting groups for amino groups of R'₁, R'₂, R'₃ and R'₄ include, for example, benzyl, benzyloxycarbonyl, t-butyloxycarbonyl and allyloxycarbonyl. Suitable protecting groups for hydroxy groups of R'₁, R'₂, R'₃ and R'₄ include, for example, benzyl, methyl, t-butyldimethylsilyl, acetyl, and tetrahydropyranyl. Suitable groups for carboxy groups of R'₁, R'₂, R'₃ and R'₄ include, but are not limited to, benzyl, allyl, methyl, ethyl and t-butyl.

Intermediate compounds of Formula II may be prepared in an enantiomeric, stereochemically homogeneous form according to the following Scheme IIA.




Formula II

Scheme IIA

or in racemic form according to Scheme IIB as described below.

Scheme IIB



5 In Scheme IIA, the starting material is an enantiomerically pure or enriched N-benzyloxycarbonyl protected α -amino acid, that may be prepared by established methods (e.g., Schotten-Baumann reaction) familiar to those skilled in the art of organic synthesis from an enantiomerically pure α -amino acid (numerous methods for preparing enantiomerically pure α -amino acids are familiar to those skilled in the art of organic synthesis, see e.g., Duthaler, *Tetrahedron* 1994, 50, 1539-1650). This starting material is coupled to a primary amino compound, R'_1NH_2 , under standard peptide bond formation conditions (e.g., DCC-HOBt), the Cbz protecting group is removed from the condensation product by catalytic hydrogenation and the resulting product is cyclized to form a hydantoin ring using CDI under non-racemizing conditions to yield a product of Formula II having either *R* or *S* absolute stereochemical configuration at C5.


15 In Scheme IIB, the starting material may either be an enantiomerically pure, enantiomerically enriched or a racemic α -N-benzyloxycarbonyl protected α -amino acid, that may be prepared by established methods (e.g., Schotten-Baumann reaction) familiar to those skilled in the art of organic synthesis as described above. This α -N-Cbz- α -amino acid starting material is coupled to a primary amino compound, R'_1NH_2 , under standard peptide bond formation conditions (DCC-HOBt), and the resulting product is cyclized directly using an excess of TBAF in THF at reflux to yield product of Formula II.

20 In the case where R'_3 is hydrogen, the product of Formula II then represents a mixture of compounds containing equal portions of C5 (*R*) and C5 (*S*) stereochemical isomers.

25 Scheme III depicts a schematic for the synthesis of the unsubstituted diphenylether sulfonyl chlorides: 4-phenoxy-phenylsulfonyl chloride. This synthesis schematic is also suitable as a general method for preparing diversely substituted diphenylether sulfonyl chlorides of Formula III to provide substituted sulfonyldiphenylether compositions

described in the invention. Compounds of Formula III may be alternatively prepared by other established procedures familiar to those skilled in the art of organic synthesis.

Scheme III



5 In cases where the primary amine $\text{R}'_1\text{NH}_2$ contains asymmetric carbon atoms, isomers of absolute *R* or *S* stereochemistry may be prepared by standard procedures resulting in compounds of Formulas IA, IB, and IC which are single isomers with homogenous stereochemistry at all asymmetric centers. Likewise, in cases where R'_2 , R'_3 , and R'_4 groups contain additional asymmetric centers, homogeneous isomers for intermediates containing these groups may be prepared by literature procedures involving
 10 either asymmetric synthesis or standard separation techniques known to one skilled in the art including resolution of chiral addition salts.

General Procedure for Preparation of Compounds of Formula IA, IB, and IC

15 To a stirred solution of Formula II at a 0.1-1.0M concentration in anhydrous THF is added 1.1 molar equivalents of LHMDS in anhydrous THF at -40°C under argon. After the addition is complete, a solution of 1.1 molar equivalents of Formula III in THF is added and the solution is allowed to warm to room temperature. After stirring for 30 min. at room temperature, saturated ammonium chloride solution is added to
 20 neutralize the solution which is then concentrated on a rotary evaporator. Ethyl acetate is added to the mixture followed by extraction with brine, separation and drying the organic phase over sodium sulfate. The organic phase is then concentrated and the resulting crude product is purified by column chromatography over silica gel. The protecting

group(s) from the so obtained product is (are) then removed by adding 5 equivalents of BBr_3 to a 0.1M solution of the product in dichloromethane at 0°C (e.g., in the case where R'_1 contains a primary amino group protected as N-Boc and R'_2 contains a phenolic hydroxy group protected as $\text{O}-\text{CH}_2\text{Ph}$). After stirring for 2 hours at room temperature, potassium bicarbonate is added to yield a neutral mixture from which the organic phase is separated, dried (sodium sulfate) and concentrated to give a crude free base. The crude free base of Formulas IA, IB, or IC is purified by column chromatography over silica gel. The purified free base is then dissolved in ethanolic HCl solution, and this solution is then concentrated to yield the corresponding hydrochloride salt of Formulas IA, IB, and IC.


General Procedure for Preparation of Enantiomerically Pure or Enriched Compounds of Formulas IA, IB, and IC

An enantiomerically pure or enriched α -N-Cbz- α -amino acid as depicted in Scheme IIA is dissolved with stirring in THF and 1.1 molar equivalents of primary amine intermediate $\text{R}'_1\text{-NH}_2$ is added in THF at room temperature. A 1:1 mixture of DCC:HOBt (1.1 molar equivalents) is then added, and the reaction is stirred until TLC or HPLC analysis indicates that the coupling reaction has proceeded to completion (typically 1-5 hour reaction time). The reaction mixture is then filtered to remove the precipitated dicyclohexylurea, concentrated *in vacuo* and purified by column chromatography over silica gel. The so obtained product is dissolved in ethanol, along with a catalytic amount a 5% palladium hydrogenation catalyst supported on activated carbon (Pd-C) and the mixture subjected to hydrogenation under an atmosphere of hydrogen. After the uptake of hydrogen by the solution is complete or TLC/HPLC analysis indicates that the reaction is complete, the mixture is filtered and concentrated to yield a crude product which is purified by column chromatography over silica gel or used directly in the subsequent step.

To a stirred solution of the hydrogenation product in THF is added a mixture of 1.1 1 molar equivalents of DIEA and 1.1 molar equivalents of CDI at -20°C . The mixture is allowed to stir at 0°C and then is heated at reflux until TLC/HPLC analysis indicates that the reaction is complete. The reaction mixture is then extracted with saturated ammonium chloride solution, organic phase is dried over sodium sulfate and concentrated to yield the crude product which is purified by column chromatography over silica gel to yield a compound of Formula II.

By way of example, a racemic mixtures of diastereomerically homogeneous compounds of Example 15 (see Table 1 below) can be prepared according to the following general procedures:

Intermediate 1



5


To a stirred solution of racemic *cis*-1,3-Bis(aminomethyl)cyclohexane (16.1 g, 74.8 mmol,) (*Helv. Chim. Acta* 1979, 62, 1065-1078) in 250mL methanol, sodium hydroxide (3.11g, 77.8 mmol,) is added at room temperature. After all the reagents are fully dissolved, di-tert-butyl dicarbonate (6.53 g, 30.0mmol,) dissolved in 80mL dioxane, is added dropwise over 1.5 hours. The mixture is stirred at room temperature for 4 hours, filtered and concentrated under reduced pressure. The resulting residue is dissolved in dichloromethane and extracted with aqueous sodium bicarbonate solution. The organic layer is dried over sodium sulfate, filtered and concentrated to give 7.25g (30.0mmol) of Intermediate 1 as colorless, viscous oil.

10


15

Data regarding the resulting Intermediate 1 are as follows: $^1\text{H-NMR}$ (CDCl_3 , 300 MHz): 0.49-0.62 (q, 1H), 0.74-0.90 (m, 2H), 1.20-1.39 (m, 3H), 1.43 (s, 9H), 1.70-1.83 (m, 4H), 2.52 (d, 2H), 2.97 (m, 2H), 3.71 (s, 2H), 4.77 (s, br, 1H).

Intermediate 2



2a



2b


20

Intermediate 1 (7.25g, 30.0mmol,) is then dissolved in 250mL dichloromethane and then alpha-carbobenzyloxy-L-tryptophan (10.2g, 30.0mmol,) and 1-hydroxybenzotriazole (7.28g, 53.9mmol) are added to the stirred solution at room temperature under nitrogen. The solution is cooled in an ice bath, and dicyclohexylcarbodiimide (11.1g, 53.9mmol,) is added. The reaction mixture is stirred at room temperature overnight, filtered and concentrated to a waxy residue. The waxy


25

residue is triturated with ether/heptane to give 12.5 g (22.2mmol) of Intermediate 2 as a white powder consisting of an equal mixture of diastereomers 2a and 2b.

Intermediate 3



3a



3b

Intermediate 2 (2.81g, 5.0mmol,) is dissolved in 55mL THF and TBAF (15mmol, 15 mL of a 1.0M solution in THF) is added. The resultant mixture is heated under reflux for 1 hour and concentrated to a residue that is dissolved in ethyl acetate and extracted with brine. The organic layer is dried over sodium sulfate, filtered and concentrated to an oil which is purified by column chromatography over silica gel (chloroform/methanol, 95:5). The fractions containing the product are evaporated and the isomers separated by repeated recrystallization from ethyl acetate/heptane (2:1) to give 1.35 g (2.97 mmol) of one diastereoisomer 3a (illustrated above) as a solid and 0.9g (1.98mmol) of the alternative diastereoisomer 3b (illustrated above) as a sticky oil. No definitive relative stereochemical assignment has been made for the respective diastereomers that are depicted in a given form for the sake of convenience.


Data regarding diastereomer 3a are as follows: MS (ESI⁺) calc. for C₂₅H₃₄N₄O₄ 454; Found 455 [M+H]⁺, ¹H-NMR (CDCl₃, 300 MHz): 0.52-0.75 (m, 2H), 0.98-1.40 (m, 5H), 1.51 (s, 9H), 1.52-1.77 (m, 4H), 1.93 (m, 1H), 2.99-3.53 (m, 3H), 4.28 (m, 2H), 4.87 (t, 1H), 6.40 (s, 1H), 7.00-7.20 (m, 3H), 7.21-7.39 (m, 1H), 7.61 (d, 1H), 10.03 (s, 1H), ¹³C-NMR (CDCl₃, 75 MHz): 24.8, 26.3, 28.4, 30.3, 33.8, 35.9, 37.3, 44.1, 47.3, 49.1, 57.5, 79.7, 107.1, 111.5, 118.5, 119.2, 121.6, 123.4, 127.7, 136.2, 156.6, 158.2, 174.1.

Data regarding the diastereomer 3b are as follows: ¹H-NMR (CDCl₃, 300 MHz): 0.50-0.72 (m, 2H), 0.83-1.30 (m, 5H), 1.48 (s, 9H), 1.50-1.80 (m, 4H), 2.38 (s, br, 1H), 2.83-3.21 (m, 3H), 4.18 (m, 2H), 4.81 (t, 1H), 6.60 (s, 1H), 7.00-7.18 (m, 3H), 7.21-7.30 (m, 1H), 7.59 (d, 1H), 9.40 (s, 1H), ¹³C-NMR (CDCl₃, 75 MHz): 24.8, 26.9, 28.3, 30.1, 34.3, 36.3, 37.6, 44.3, 46.8, 57.9, 79.3, 107.9, 111.3, 118.5, 119.2, 121.7, 123.7, 127.2, 136.1, 156.3, 157.9, 174.0.

To a stirred solution of 4-benzyloxy-3-chlorophenol (4.69g, 20mmol) in 120mL DMF, potassium carbonate (4.14 g, 30mmol), 2-chloro-5-nitro-N,N-dimethylbenzamide (4.57g, 20mmol), and 18-crown-6 (0.2g, 0.75mmol) are added at room temperature. The mixture is heated at 100°C for 18 hours, cooled to room temperature, filtered, and concentrated. The resulting residue is dissolved in ethyl acetate and washed with aqueous sodium bicarbonate. The organic layer is concentrated, dissolved in ethanol, stirred with activated charcoal for 15 minutes, filtered, and the solvent evaporated to give 8.2g (19.2 mmol) of the diphenylether coupling product to be employed for the following reduction step without further purification.

Specifically, the above obtained diphenylether coupling product is dissolved in 100mL acetic acid/water (1:1) and iron powder (10.76 g, 192mmol) was added to that solution. The mixture is heated at reflux for 1 hour, cooled to room temperature and 15% sodium hydroxide solution added to adjust the solution to neutral pH. The mixture is filtered and the solid is washed with ethyl acetate. The combined filtrates are washed with water and the organic layer is concentrated and purified by column chromatography over silica gel (ethyl acetate) to give 5.48 g (13.8mmol) of aniline product as a pale yellow solid.


Intermediate 4



The above obtained 5.48g of aniline is then dissolved in 50mL acetic acid containing 5mL concentrated hydrochloric acid and cooled at -5°C. Sodium nitrite (1.14 g, 16.6mmol) dissolved in 5mL water is added dropwise followed by stirring at -5°C for 1 hour to yield a diazonium salt solution. A separate solution is prepared by adding CuCl₂ dihydrate (472mg, 2.76mmol) in 1 ml water to 50mL of SO₂-saturated acetic acid. The diazonium salt solution is added to this solution at 0°C with stirring, and the solution allowed to stand overnight at 4°C. The resulting light green solution is poured into 500mL ice water and extracted with dichloromethane. The combined organic layers are washed with water, dried over sodium sulfate, and concentrated. The obtained residue is purified by column chromatography over silica gel (ethyl acetate/heptane 2:1) to give 4.71g (9.82 mmol) of Intermediate 4 as a yellow foam.

Data regarding Intermediate 4 are as follows: MS (ESI⁺) calc. for C₂₂H₁₉Cl₂NO₅S 480; Found 481 [M+H]⁺, ¹H-NMR (CDCl₃, 300MHz): 2.98 (s, 3H), 3.11 (s, 3H), 5.19 (s, 2H), 6.86-7.05 (m, 3H), 7.28 (d, 1H), 7.30-7.52 (m, 5H), 7.92 (dd, 1H), 8.04 (d, 1H), ¹³C-NMR (CDCl₃, 75MHz): 34.9, 38.3, 71.2, 114.9, 115.8, 119.8, 122.9, 124.4, 127.0, 128.1, 128.2, 128.6, 129.7, 135.9, 137.9, 147.0, 152.2, 159.3, 165.5.

Intermediate 5a




To a stirred solution of Intermediate 3a (454 mg, 1.0 mmol) in 10mL anhydrous THF, LHMDS (0.9mmol, 0.9mL of a 1M solution in hexane) is added at -10°C under nitrogen. The mixture is stirred at this temperature for 20 minutes, followed by the addition of Intermediate 4 (570mg, 1.2mmol) in 5mL THF. The reaction mixture is allowed to warm to room temperature and is stirred for 1.5 hours. The reaction is quenched by addition of aqueous ammonium chloride solution, concentrated, and

partitioned between brine and ethyl acetate. The organic phase was separated, dried over sodium sulfate, filtered, and concentrated to a viscous oil. The oil is purified by column chromatography over silica gel (ethyl acetate/heptane) to give 575 mg (0.64 mmol) of Intermediate 5a as a white foam.

5 Data regarding Intermediate 5a are as follows: $^1\text{H-NMR}$ (CDCl_3 , 300 MHz): 0.20-0.59 (m, 2H), 0.62-1.18 (m, 4H), 1.53 (s, 9H), 1.60 (m, 2H), 2.75 (t, 1H), 2.98 (s, 3H), 3.10 (s, 3H), 3.10 (t, 1H), 3.71 (m, 2H), 4.74 (s, 1H), 4.93 (t, 1H), 5.09 (s, 2H), 6.80 (d, 1H), 6.84-7.50 (m, 12H), 7.69 (m, 1H), 8.01 (m, 2H), 10.06 (s, 1H).

Intermediate 5b




10

Substituting Intermediate 3b for Intermediate 3a and otherwise following the same procedure used to prepare Intermediate 5a, 396 mg (0.44 mmol) of Intermediate 5b were obtained as a white foam.


15 Data regarding Intermediate 5b are as follows: $^1\text{H-NMR}$ (CDCl_3 , 300 MHz): 0.05-1.35 (m, 6H), 1.48 (m, 4H), 1.68 (s, 9H), 1.70 (m, 2H), 2.90 (m, 1H), 3.09 (m, 1H), 3.14 (s, 3H), 3.30 (s, 3H), 3.30 (m, 1H), 3.91 (m, 2H), 4.82 (t, 1H), 4.93 (s, 1H), 5.39 (s, 2H), 7.01 (d, 1H), 7.09-7.69 (m, 12H), 7.88 (m, 1H), 8.21 (m, 2H), 9.44 (s, 1H).

37

Intermediate 6a

A mixture of Intermediate 5a (494mg, 0.55mmol), 49mg of 10% palladium on carbon, and ammonium formate (346mg, 5.5mmol,) in 10mL of isopropyl acetate/isopropanol (1:1vvl) containing 50microliters of concentrated HCl is heated at 85°C for 1 hour. It is then filtered, the filtrate evaporated, and the residue purified by column chromatography (ethyl acetate/heptane 2:1) to give 399 mg (0.494 mmol, 90 %) of Intermediate 6a as a white foam.

Data regarding Intermediate 6a are as follows: ¹H-NMR (CDCl₃, 300 MHz): 0.20-0.59 (m, 3H), 0.62-0.92 (m, 3H), 1.53 (s, 9H), 1.60 (m, 2H), 2.72 (t, 1H), 2.99 (s, 3H), 3.10 (m, 1H), 3.19 (s, 3H), 3.71 (m, 2H), 4.72 (s, 1H), 4.90 (t, 1H), 6.81 (d, 1H), 6.84-7.28 (m, 7H), 7.67 (m, 1H), 8.03 (m, 2H), 10.02 (s, 1H).


Intermediate 6b:

A mixture of Intermediate 5b (670mg, 0.75mmol), 67mg of 10% palladium on carbon, and ammonium formate (470 mg, 7.5mmol,) in 10mL of isopropyl acetate/isopropanol (1:1/vv) containing 60microliters of concentrated HCl is heated at 85°C for 1 hour. It is then filtered, the filtrate evaporated, and the residue purified by

column chromatography (ethyl acetate/heptane 2:1) to give 505 mg (0.63 mmol) of Intermediate 6b as a white foam.

Data regarding Intermediate 6b are as follows: $^1\text{H-NMR}$ (CDCl_3 , 300 MHz): 0.06-1.28 (m, 8H), 1.49 (s, 9H), 1.52 (m, 4H), 2.69 (m, 1H), 2.89 (m, 1H), 2.99 (s, 3H), 3.12 (m, 1H), 3.27 (s, 3H), 3.69 (m, 2H), 4.63 (t, 1H), 4.78 (s, 1H), 6.79-7.30 (m, 8H), 7.67 (m, 1H), 8.01 (m, 2H), 9.22 (s, 1H).

Diastereomer of Example 15: Example 15a




Intermediate 6a (290mg, 0.36mmol,) is dissolved in 2 ml of 50% trifluoroacetic acid in dichloromethane. The resulting solution is stirred at room temperature for 1 hour. Sodium bicarbonate is added to adjust the solution to pH 8, and the mixture is partitioned between ethyl acetate and aqueous sodium bicarbonate. The organic layer is dried over sodium sulfate, filtered, and concentrated to a viscous oil. The resulting oil is purified by column chromatography over silica gel (chloroform/methanol/ammonium hydroxide) to give 148mg (0.21mmol) of a diastereomer of Compound Example 15 (hereinafter Example 15a, see Table 1 below) in free base form as a white powder. The corresponding hydrochloride salt was prepared by dissolving the free base (148mg, 0.21 mmol,) in 2mL 0.2M ethanolic HCl solution, heating the solution at 35°C for 10 minutes, and then evaporating the solvent. The resulting solid is washed with ether and dried in vacuum to give 126 mg (0.17mmol) of the hydrochloride salt as a light pink powder.

Data regarding Compound Example 15a are as follows: MS (ESI^+) calc. for $\text{C}_{35}\text{H}_{38}\text{ClN}_5\text{O}_7\text{S}$ 707; Found 708 $[\text{M}+\text{H}]^+$, $^1\text{H-NMR}$ (CD_3OD , 300 MHz): 0.03-0.13 (m, 1H), 0.16-0.34 (m, 1H), 0.47-0.64 (m, 1H), 0.65-0.76 (m, 1H), 0.83-1.11 (m, 2H), 1.50 (m, 2H), 2.40 (m, 2H), 2.87 (m, 1H), 2.98 (s, 3H), 3.06 (m, 1H), 3.10 (s, 3H), 3.25 (m,

2H), 3.55-3.76 (m, 2H), 4.98 (m, 1H), 6.90-7.17 (m, 7H), 7.32 (d, 1H), 7.61 (d, 1H), 8.09 (m, 2H).

Diastereomer of Example 15: Example 15b



5 The free base and hydrochloride salt (120mg, 0.16mmol) of an alternate diastereomer of Compound Example 15 (hereinafter Example 15b; see Table 1 below) was prepared in analogous fashion to Example 15a above by substituting Intermediate 6b (0.557mmol, 450mg) for Intermediate 6a.

10 Data regarding Compound Example 15b are as follows: MS (ESI⁺) calc. for C₃₅H₃₈ClN₅O₇S 707; Found 708 [M+H]⁺, ¹H-NMR (CD₃OD, 300 MHz): 0.10-0.28 (m, 1H), 0.44 (m, 1H), 0.59 (m, 1H), 0.80 (m, 1H), 3.5 (m, 1H), 1.52 (m, 1H), 2.49 (m, 2H), 2.84 (m, 2H), 2.96 (s, 3H), 3.05 (m, 1H), 3.11 (s, 3H), 3.28 (m, 2H), 3.55-3.74 (m, 2H), 4.97 (m, 1H), 6.90-7.10 (m, 6H), 7.16 (m, 1H), 7.30 (d, 1H), 7.60 (d, 1H), 8.08 (m, 2H).

15 Figure 1 illustrates the logarithmic concentration-response curves for inhibition of [¹²⁵I]Somatostatin binding to human SSTR2 receptors expressed in CHO-K1 cells by somatostatin as compared to binding by Compound Example 15a (see Table I) hydrochloride salt. Test compounds were tested in duplicate at all concentrations. The average percent inhibition of duplicate was used for plotting the displacement curves, fitted by non-linear regression. IC₅₀ is the concentration of the test substance that causes
20 50% displacement of [¹²⁵I] somatostatin binding from SSTR2 receptor binding sites. The following Chart I provides the IC₅₀ data for both Example 15a HCl salt and somatostatin.

Chart I

Compound	IC ₅₀
● Example 15a HCl salt	0.290 ±0.0395 nM
■ Somatostatin	3.69±0.146 pM

Figure 2 illustrates the logarithmic concentration-response curves for inhibition of [¹²⁵I]Somatostatin binding to human SSTR2 receptors expressed in CHO-K1 cells by somatostatin and Compound Example 15b (see Table I) hydrochloride salt. Test compounds were tested in duplicate at all concentrations. The average percent inhibition of duplicate was used for plotting the displacement curves, fitted by non-linear regression. IC₅₀ is the concentration of the test substance that causes 50% displacement of [¹²⁵I] somatostatin binding from somatostatin SSTR2 receptor binding sites. The following Chart II provides the IC₅₀ data for both Example 15a HCl salt and somatostatin.

Chart II

Compound	IC ₅₀
● Example 15b HCl salt	1.30 ±0.183 nM
■ Somatostatin	3.69±0.147 pM

Examples of compounds of the invention include those defined in the following tables and are prepared in analogous manner to procedures defined above.




Table 1

Example	a	b	c
1	OH	H	H
2	NH ₂	H	H
3	F	H	H
4	CF ₃	H	H
5	OH	Cl	H
6	NH ₂	Cl	H
7	F	Cl	H
8	Cl	Cl	H
9	OH	H	C(O)N(CH ₃) ₂
10	NH ₂	H	C(O)N(CH ₃) ₂
11	F	H	C(O)N(CH ₃) ₂
12	Cl	H	C(O)N(CH ₃) ₂
13	OCH ₃	H	C(O)N(CH ₃) ₂
14	CF ₃	H	C(O)N(CH ₃) ₂
15	OH	Cl	C(O)N(CH ₃) ₂
16	NH ₂	Cl	C(O)N(CH ₃) ₂
17	Cl	Cl	C(O)N(CH ₃) ₂
18	H	Cl	C(O)N(CH ₃) ₂
19	OCH ₃	Cl	C(O)N(CH ₃) ₂
20	CF ₃	Cl	C(O)N(CH ₃) ₂
21	OH	H	C(O)N(CH ₂ CH ₂ O) ₂ =morpholine amido
22	NH ₂	H	C(O)N(CH ₂ CH ₂ O) ₂
23	F	H	C(O)N(CH ₂ CH ₂ O) ₂
24	Cl	H	C(O)N(CH ₂ CH ₂ O) ₂

Example	a	b	c
25	OCH ₃	H	C(O)N(CH ₂ CH ₂ O) ₂
26	CF ₃	H	C(O)N(CH ₂ CH ₂ O) ₂
27	OH	Cl	C(O)N(CH ₂ CH ₂ O) ₂
28	NH ₂	Cl	C(O)N(CH ₂ CH ₂ O) ₂
29	Cl	Cl	C(O)N(CH ₂ CH ₂ O) ₂
30	OCH ₃	Cl	C(O)N(CH ₂ CH ₂ O) ₂
31	OH	H	CN = cyano
32	NH ₂	H	CN
33	F	H	CN
34	Cl	H	CN
35	OCH ₃	H	CN
36	CF ₃	H	CN
37	OH	Cl	CN
38	NH ₂	Cl	CN
39	Cl	Cl	CN
40	OCH ₃	Cl	CN
41	OH	H	Cl
42	NH ₂	H	Cl
43	F	H	Cl
44	Cl	H	Cl
45	OCH ₃	H	Cl
46	CF ₃	H	Cl
47	OH	Cl	Cl
48	NH ₂	Cl	Cl
49	Cl	Cl	Cl
50	OCH ₃	Cl	Cl




Table 2

Example	a	b	c
51	OH	H	H
52	NH ₂	H	H
53	F	H	H
54	CF ₃	H	H
55	OH	Cl	H
56	NH ₂	Cl	H
57	F	Cl	H
58	Cl	Cl	H
59	OH	H	C(O)N(CH ₃) ₂
60	NH ₂	H	C(O)N(CH ₃) ₂
61	F	H	C(O)N(CH ₃) ₂
62	Cl	H	C(O)N(CH ₃) ₂
63	OCH ₃	H	C(O)N(CH ₃) ₂
64	CF ₃	H	C(O)N(CH ₃) ₂
65	OH	Cl	C(O)N(CH ₃) ₂
66	NH ₂	Cl	C(O)N(CH ₃) ₂
67	Cl	Cl	C(O)N(CH ₃) ₂
68	H	Cl	C(O)N(CH ₃) ₂
69	OCH ₃	Cl	C(O)N(CH ₃) ₂
70	CF ₃	Cl	C(O)N(CH ₃) ₂
71	OH	H	C(O)N(CH ₂ CH ₂ O) ₂ =morpholine amido
72	NH ₂	H	C(O)N(CH ₂ CH ₂ O) ₂
73	F	H	C(O)N(CH ₂ CH ₂ O) ₂
74	Cl	H	C(O)N(CH ₂ CH ₂ O) ₂

Example	a	b	c
75	OCH ₃	H	C(O)N(CH ₂ CH ₂ O) ₂
76	CF ₃	H	C(O)N(CH ₂ CH ₂ O) ₂
77	OH	Cl	C(O)N(CH ₂ CH ₂ O) ₂
78	NH ₂	Cl	C(O)N(CH ₂ CH ₂ O) ₂
79	Cl	Cl	C(O)N(CH ₂ CH ₂ O) ₂
80	OCH ₃	Cl	C(O)N(CH ₂ CH ₂ O) ₂




Table 3

Example	a	b	c	d
81	OH	H	H	CO ₂ CH ₃
82	NH ₂	H	H	CO ₂ CH ₃
83	OH	Cl	H	CO ₂ CH ₃
84	OH	H	C(O)N(CH ₃) ₂	CO ₂ CH ₃
85	NH ₂	H	C(O)N(CH ₃) ₂	CO ₂ CH ₃
86	F	H	C(O)N(CH ₃) ₂	CO ₂ CH ₃
87	OH	Cl	C(O)N(CH ₃) ₂	CO ₂ CH ₃
88	OH	H	C(O)N(CH ₂ CH ₂ O) ₂ = morpholine amido	CO ₂ CH ₃
89	NH ₂	H	C(O)N(CH ₂ CH ₂ O) ₂	CO ₂ CH ₃
90	F	H	C(O)N(CH ₂ CH ₂ O) ₂	CO ₂ CH ₃
91	OH	Cl	C(O)N(CH ₂ CH ₂ O) ₂	CO ₂ CH ₃
92	OH	H	CN = cyano	CO ₂ CH ₃
93	NH ₂	H	CN	CO ₂ CH ₃
94	F	H	CN	CO ₂ CH ₃
95	OH	Cl	CN	CO ₂ CH ₃
96	OH	H	Cl	CO ₂ CH ₃
97	NH ₂	H	Cl	CO ₂ CH ₃
98	F	H	Cl	CO ₂ CH ₃
99	OH	Cl	Cl	CO ₂ CH ₃
100	OH	H	H	CO ₂ CH ₂ CH ₃
101	NH ₂	H	H	CO ₂ CH ₂ CH ₃
102	OH	Cl	H	CO ₂ CH ₂ CH ₃
103	OH	H	C(O)N(CH ₃) ₂	CO ₂ CH ₂ CH ₃
104	NH ₂	H	C(O)N(CH ₃) ₂	CO ₂ CH ₂ CH ₃
105	F	H	C(O)N(CH ₃) ₂	CO ₂ CH ₂ CH ₃

Example	a	b	c	d
106	OH	Cl	$C(O)N(CH_3)_2$	$CO_2CH_2CH_3$
107	OH	H	$C(O)N(CH_2CH_2O)_2$ =morpholine amido	$CO_2CH_2CH_3$
108	NH ₂	H	$C(O)N(CH_2CH_2O)_2$	$CO_2CH_2CH_3$
109	F	H	$C(O)N(CH_2CH_2O)_2$	$CO_2CH_2CH_3$
110	OH	Cl	$C(O)N(CH_2CH_2O)_2$	$CO_2CH_2CH_3$
111	OH	H	CN = cyano	$CO_2CH_2CH_3$
112	NH ₂	H	CN	$CO_2CH_2CH_3$
113	F	H	CN	$CO_2CH_2CH_3$
114	OH	Cl	CN	$CO_2CH_2CH_3$
115	OH	H	Cl	$CO_2CH_2CH_3$
116	NH ₂	H	Cl	$CO_2CH_2CH_3$
117	F	H	Cl	$CO_2CH_2CH_3$
118	OH	Cl	Cl	$CO_2CH_2CH_3$
119	OH	H	H	CO_2 -tert-butyl
120	NH ₂	H	H	CO_2 -tert-butyl
121	OH	Cl	H	CO_2 -tert-butyl
122	OH	H	$C(O)N(CH_3)_2$	CO_2 -tert-butyl
123	NH ₂	H	$C(O)N(CH_3)_2$	CO_2 -tert-butyl
124	F	H	$C(O)N(CH_3)_2$	CO_2 -tert-butyl
125	OH	Cl	$C(O)N(CH_3)_2$	CO_2 -tert-butyl
126	OH	H	$C(O)N(CH_2CH_2O)_2$ =morpholine amido	CO_2 -tert-butyl
127	NH ₂	H	$C(O)N(CH_2CH_2O)_2$	CO_2 -tert-butyl
128	F	H	$C(O)N(CH_2CH_2O)_2$	CO_2 -tert-butyl
129	OH	Cl	$C(O)N(CH_2CH_2O)_2$	CO_2 -tert-butyl
130	OH	H	CN = cyano	CO_2 -tert-butyl
131	NH ₂	H	CN	CO_2 -tert-butyl
132	F	H	CN	CO_2 -tert-butyl
133	OH	Cl	CN	CO_2 -tert-butyl
134	OH	H	Cl	CO_2 -tert-butyl
135	NH ₂	H	Cl	CO_2 -tert-butyl
136	F	H	Cl	CO_2 -tert-butyl
137	OH	Cl	Cl	CO_2 -tert-butyl
138	OH	H	H	phenyl
139	NH ₂	H	H	phenyl
140	OH	Cl	H	phenyl

Example	a	b	c	d
141	OH	H	$C(O)N(CH_3)_2$	phenyl
142	NH ₂	H	$C(O)N(CH_3)_2$	phenyl
143	F	H	$C(O)N(CH_3)_2$	phenyl
144	OH	Cl	$C(O)N(CH_3)_2$	phenyl
145	OH	H	$C(O)N(CH_2CH_2O)_2$ =morpholine amido	phenyl
146	NH ₂	H	$C(O)N(CH_2CH_2O)_2$	phenyl
147	F	H	$C(O)N(CH_2CH_2O)_2$	phenyl
148	OH	Cl	$C(O)N(CH_2CH_2O)_2$	phenyl
149	OH	H	CN = cyano	phenyl
150	NH ₂	H	CN	phenyl
151	F	H	CN	phenyl
152	OH	Cl	CN	phenyl
153	OH	H	Cl	phenyl
154	NH ₂	H	Cl	phenyl
155	F	H	Cl	phenyl
156	OH	Cl	Cl	phenyl
157	OH	H	H	CN =cyano
158	NH ₂	H	H	CN =cyano
159	OH	Cl	H	CN =cyano
160	OH	H	$C(O)N(CH_3)_2$	CN =cyano
161	NH ₂	H	$C(O)N(CH_3)_2$	CN =cyano
162	F	H	$C(O)N(CH_3)_2$	CN =cyano
163	OH	Cl	$C(O)N(CH_3)_2$	CN =cyano
164	OH	H	$C(O)N(CH_2CH_2O)_2$ =morpholine amido	CN =cyano
165	NH ₂	H	$C(O)N(CH_2CH_2O)_2$	CN =cyano
166	F	H	$C(O)N(CH_2CH_2O)_2$	CN =cyano
167	OH	Cl	$C(O)N(CH_2CH_2O)_2$	CN =cyano
168	OH	H	CN = cyano	CN =cyano
169	NH ₂	H	CN	CN =cyano
170	F	H	CN	CN =cyano
171	OH	Cl	CN	CN =cyano
172	OH	H	Cl	CN =cyano
173	NH ₂	H	Cl	CN =cyano
174	F	H	Cl	CN =cyano
175	OH	Cl	Cl	CN =cyano




Table 4

Example	a	b	c
176	OH	H	H
177	NH ₂	H	H
178	OH	Cl	H
179	OH	H	C(O)N(CH ₃) ₂
180	NH ₂	H	C(O)N(CH ₃) ₂
181	F	H	C(O)N(CH ₃) ₂
182	OH	Cl	C(O)N(CH ₃) ₂
183	OH	H	C(O)N(CH ₂ CH ₂ O) ₂ =morpholine amido
184	NH ₂	H	C(O)N(CH ₂ CH ₂ O) ₂
185	F	H	C(O)N(CH ₂ CH ₂ O) ₂
186	OH	Cl	C(O)N(CH ₂ CH ₂ O) ₂
187	OH	H	CN = cyano
188	NH ₂	H	CN
189	F	H	CN
190	OH	Cl	CN
191	OH	H	Cl
192	NH ₂	H	Cl
193	F	H	Cl
194	OH	Cl	Cl




Table 5

Example	a	b	C	d
195	OH	H	H	CO ₂ CH ₃
196	NH ₂	H	H	CO ₂ CH ₃
197	OH	Cl	H	CO ₂ CH ₃
198	OH	H	C(O)N(CH ₃) ₂	CO ₂ CH ₃
199	NH ₂	H	C(O)N(CH ₃) ₂	CO ₂ CH ₃
200	F	H	C(O)N(CH ₃) ₂	CO ₂ CH ₃
201	OH	Cl	C(O)N(CH ₃) ₂	CO ₂ CH ₃
202	OH	H	C(O)N(CH ₂ CH ₂ O) ₂ =morpholine amido	CO ₂ CH ₃
203	NH ₂	H	C(O)N(CH ₂ CH ₂ O) ₂	CO ₂ CH ₃
204	F	H	C(O)N(CH ₂ CH ₂ O) ₂	CO ₂ CH ₃
205	OH	Cl	C(O)N(CH ₂ CH ₂ O) ₂	CO ₂ CH ₃
206	OH	H	CN = cyano	CO ₂ CH ₃
207	NH ₂	H	CN	CO ₂ CH ₃
208	F	H	CN	CO ₂ CH ₃
209	OH	Cl	CN	CO ₂ CH ₃
210	OH	H	Cl	CO ₂ CH ₃
211	NH ₂	H	Cl	CO ₂ CH ₃
212	F	H	Cl	CO ₂ CH ₃
213	OH	Cl	Cl	CO ₂ CH ₃
214	OH	H	H	CO ₂ CH ₂ CH ₃
215	NH ₂	H	H	CO ₂ CH ₂ CH ₃
216	OH	Cl	H	CO ₂ CH ₂ CH ₃
217	OH	H	C(O)N(CH ₃) ₂	CO ₂ CH ₂ CH ₃
218	NH ₂	H	C(O)N(CH ₃) ₂	CO ₂ CH ₂ CH ₃

Example	a	b	C	d
219	F	H	$C(O)N(CH_3)_2$	$CO_2CH_2CH_3$
220	OH	Cl	$C(O)N(CH_3)_2$	$CO_2CH_2CH_3$
221	OH	H	$C(O)N(CH_2CH_2O)_2$ =morpholine amido	$CO_2CH_2CH_3$
222	NH ₂	H	$C(O)N(CH_2CH_2O)_2$	$CO_2CH_2CH_3$
223	F	H	$C(O)N(CH_2CH_2O)_2$	$CO_2CH_2CH_3$
224	OH	Cl	$C(O)N(CH_2CH_2O)_2$	$CO_2CH_2CH_3$
225	OH	H	CN = cyano	$CO_2CH_2CH_3$
226	NH ₂	H	CN	$CO_2CH_2CH_3$
227	F	H	CN	$CO_2CH_2CH_3$
228	OH	Cl	CN	$CO_2CH_2CH_3$
229	OH	H	Cl	$CO_2CH_2CH_3$
230	NH ₂	H	Cl	$CO_2CH_2CH_3$
231	F	H	Cl	$CO_2CH_2CH_3$
232	OH	Cl	Cl	$CO_2CH_2CH_3$
233	OH	H	H	CO_2 -tert-butyl
234	NH ₂	H	H	CO_2 -tert-butyl
235	OH	Cl	H	CO_2 -tert-butyl
236	OH	H	$C(O)N(CH_3)_2$	CO_2 -tert-butyl
237	NH ₂	H	$C(O)N(CH_3)_2$	CO_2 -tert-butyl
238	F	H	$C(O)N(CH_3)_2$	CO_2 -tert-butyl
239	OH	Cl	$C(O)N(CH_3)_2$	CO_2 -tert-butyl
240	OH	H	$C(O)N(CH_2CH_2O)_2$ =morpholine amido	CO_2 -tert-butyl
241	NH ₂	H	$C(O)N(CH_2CH_2O)_2$	CO_2 -tert-butyl
242	F	H	$C(O)N(CH_2CH_2O)_2$	CO_2 -tert-butyl
243	OH	Cl	$C(O)N(CH_2CH_2O)_2$	CO_2 -tert-butyl
244	OH	H	CN = cyano	CO_2 -tert-butyl
245	NH ₂	H	CN	CO_2 -tert-butyl
246	F	H	CN	CO_2 -tert-butyl
247	OH	Cl	CN	CO_2 -tert-butyl
248	OH	H	Cl	CO_2 -tert-butyl
249	NH ₂	H	Cl	CO_2 -tert-butyl
250	F	H	Cl	CO_2 -tert-butyl
251	OH	Cl	Cl	CO_2 -tert-butyl




Table 6

Example	a	b	c
252	OH	H	H
253	NH ₂	H	H
254	F	H	H
255	CF ₃	H	H
256	OH	Cl	H
257	NH ₂	Cl	H
258	F	Cl	H
259	Cl	Cl	H
260	OH	H	C(O)N(CH ₃) ₂
261	NH ₂	H	C(O)N(CH ₃) ₂
262	F	H	C(O)N(CH ₃) ₂
263	Cl	H	C(O)N(CH ₃) ₂
264	OCH ₃	H	C(O)N(CH ₃) ₂
265	CF ₃	H	C(O)N(CH ₃) ₂
266	OH	Cl	C(O)N(CH ₃) ₂
267	NH ₂	Cl	C(O)N(CH ₃) ₂
268	Cl	Cl	C(O)N(CH ₃) ₂
269	H	Cl	C(O)N(CH ₃) ₂
270	OCH ₃	Cl	C(O)N(CH ₃) ₂
271	CF ₃	Cl	C(O)N(CH ₃) ₂
272	OH	H	C(O)N(CH ₂ CH ₂ O) ₂ =morpholine amido
273	NH ₂	H	C(O)N(CH ₂ CH ₂ O) ₂
274	F	H	C(O)N(CH ₂ CH ₂ O) ₂

Example	a	b	c
275	Cl	H	$C(O)N(CH_2CH_2O)_2$
276	OCH ₃	H	$C(O)N(CH_2CH_2O)_2$
277	CF ₃	H	$C(O)N(CH_2CH_2O)_2$
278	OH	Cl	$C(O)N(CH_2CH_2O)_2$
279	NH ₂	Cl	$C(O)N(CH_2CH_2O)_2$
280	Cl	Cl	$C(O)N(CH_2CH_2O)_2$
281	OCH ₃	Cl	$C(O)N(CH_2CH_2O)_2$
282	OH	H	CN = cyano
283	NH ₂	H	CN
284	F	H	CN
285	Cl	H	CN
286	OCH ₃	H	CN
287	CF ₃	H	CN
288	OH	Cl	CN
289	NH ₂	Cl	CN
290	Cl	Cl	CN
291	OCH ₃	Cl	CN
292	OH	H	Cl
293	NH ₂	H	Cl
294	F	H	Cl
295	Cl	H	Cl
296	OCH ₃	H	Cl
297	CF ₃	H	Cl
298	OH	Cl	Cl
299	NH ₂	Cl	Cl
300	Cl	Cl	Cl
301	OCH ₃	Cl	Cl

53




Table 7

Example	a	b	c
302	OH	H	H
303	NH ₂	H	H
304	F	H	H
305	CF ₃	H	H
306	OH	Cl	H
307	NH ₂	Cl	H
308	F	Cl	H
309	Cl	Cl	H
310	OH	H	C(O)N(CH ₃) ₂
311	NH ₂	H	C(O)N(CH ₃) ₂
312	F	H	C(O)N(CH ₃) ₂
313	Cl	H	C(O)N(CH ₃) ₂
314	OCH ₃	H	C(O)N(CH ₃) ₂
315	CF ₃	H	C(O)N(CH ₃) ₂
316	OH	Cl	C(O)N(CH ₃) ₂
317	NH ₂	Cl	C(O)N(CH ₃) ₂
318	Cl	Cl	C(O)N(CH ₃) ₂
319	H	Cl	C(O)N(CH ₃) ₂
320	OCH ₃	Cl	C(O)N(CH ₃) ₂
321	CF ₃	Cl	C(O)N(CH ₃) ₂
322	OH	H	C(O)N(CH ₂ CH ₂ O) ₂ =morpholine amido
323	NH ₂	H	C(O)N(CH ₂ CH ₂ O) ₂
324	F	H	C(O)N(CH ₂ CH ₂ O) ₂

Example	a	b	C
325	Cl	H	$C(O)N(CH_2CH_2O)_2$
326	OCH_3	H	$\dot{C}(O)N(CH_2CH_2O)_2$
327	CF_3	H	$C(O)N(CH_2CH_2O)_2$
328	OH	Cl	$C(O)N(CH_2CH_2O)_2$
329	NH_2	Cl	$C(O)N(CH_2CH_2O)_2$
330	Cl	Cl	$C(O)N(CH_2CH_2O)_2$
331	OCH_3	Cl	$C(O)N(CH_2CH_2O)_2$
332	OH	H	CN = cyano
333	NH_2	H	CN
334	F	H	CN
335	Cl	H	CN
336	OCH_3	H	CN
337	CF_3	H	CN
338	OH	Cl	CN
339	NH_2	Cl	CN
340	Cl	Cl	CN
341	OCH_3	Cl	CN
342	OH	H	Cl
343	NH_2	H	Cl
344	F	H	Cl
345	Cl	H	Cl
346	OCH_3	H	Cl
347	CF_3	H	Cl
348	OH	Cl	Cl
349	NH_2	Cl	Cl
350	Cl	Cl	Cl
351	OCH_3	Cl	Cl




Table 8

Example	a	b	c
352	OH	H	H
353	NH ₂	H	H
354	F	H	H
355	CF ₃	H	H
356	OH	Cl	H
357	NH ₂	Cl	H
358	F	Cl	H
359	Cl	Cl	H
360	OH	H	C(O)N(CH ₃) ₂
361	NH ₂	H	C(O)N(CH ₃) ₂
362	F	H	C(O)N(CH ₃) ₂
363	Cl	H	C(O)N(CH ₃) ₂
364	OCH ₃	H	C(O)N(CH ₃) ₂
365	CF ₃	H	C(O)N(CH ₃) ₂
366	OH	Cl	C(O)N(CH ₃) ₂
367	NH ₂	Cl	C(O)N(CH ₃) ₂
368	Cl	Cl	C(O)N(CH ₃) ₂
369	H	Cl	C(O)N(CH ₃) ₂
370	OCH ₃	Cl	C(O)N(CH ₃) ₂
371	CF ₃	Cl	C(O)N(CH ₃) ₂
372	OH	H	C(O)N(CH ₂ CH ₂ O) ₂ =morpholine amido


Example	a	b	c
373	NH ₂	H	C(O)N(CH ₂ CH ₂ O) ₂
374	F	H	Ċ(O)N(CH ₂ CH ₂ O) ₂
375	Cl	H	C(O)N(CH ₂ CH ₂ O) ₂
376	OCH ₃	H	C(O)N(CH ₂ CH ₂ O) ₂
377	CF ₃	H	C(O)N(CH ₂ CH ₂ O) ₂
378	OH	Cl	C(O)N(CH ₂ CH ₂ O) ₂
379	NH ₂	Cl	C(O)N(CH ₂ CH ₂ O) ₂
380	Cl	Cl	C(O)N(CH ₂ CH ₂ O) ₂
381	OCH ₃	Cl	C(O)N(CH ₂ CH ₂ O) ₂
382	OH	H	CN = cyano
383	NH ₂	H	CN
384	F	H	CN
385	Cl	H	CN
386	OCH ₃	H	CN
387	CF ₃	H	CN
388	OH	Cl	CN
389	NH ₂	Cl	CN
390	Cl	Cl	CN
391	OCH ₃	Cl	CN
392	OH	H	Cl
393	NH ₂	H	Cl
394	F	H	Cl
395	Cl	H	Cl
396	OCH ₃	H	Cl
397	CF ₃	H	Cl
398	OH	Cl	Cl
399	NH ₂	Cl	Cl
400	Cl	Cl	Cl
401	OCH ₃	Cl	Cl

It should be understood that the examples and embodiments described herein are for illustrative purposes only and that various modifications or changes in light thereof will be suggested to persons skilled in the art and are to be included within the spirit and purview of this application and the scope of the appended claims.

Claims

We claim:

1. A non-peptide, somatostatin receptor ligand represented by Formula IC:




Formula IC



or a pharmaceutically acceptable racemate, enantiomer, diastereomer, solvate, salt or prodrug thereof, wherein:

A and B independently are oxygen, or (H,H);

R₁ is a group selected from Formula Set A or Formula Set B



Formula Set A



Formula Set B

wherein;

each R_a independently is hydrogen, alkyl, alkenyl, alkynyl, hydroxyalkyl, alkoxyalkyl, arylalkyl, or form together a 3-6 carbon-membered ring;

G is aryloxy, cyano, $-\text{CO}_2\text{R}_a$, or $-\text{C}(\text{O})\text{N}(\text{R}_b)_2$ wherein R_b is independently hydrogen, alkyl, alkenyl, alkynyl, hydroxyalkyl, alkoxyalkyl, monoalkylaminoalkyl, dialkylaminoalkyl, aminoalkyl, arylalkyl, heterocyclalkyl, or is optionally joined to a five- or six-membered ring;

Q is a C_1 - C_5 alkyl, alkenyl, or alkynyl linker group that can optionally include oxygen, sulfur, or nitrogen atoms or $-\text{N}(\text{R}_a)-$ and/or is optionally substituted by hydroxy, alkoxy, alkyl, or aryl;

each E_1 , E_2 , and E_3 independently is a covalent bond, carbon, oxygen, sulfur, or nitrogen optionally substituted with an alkyl, alkenyl, alkynyl, alkylaryl, or aryl, n is an independently 0, 1, or 2; and


J is hydrogen, alkyl, heterocyclalkyl, or $\text{CH}_2\text{N}(\text{R}_a)_2$;

R_2 is $-\text{SO}_2\text{-Ar}$ or $(\text{CHR}_c)_n\text{-Ar}$

wherein

R_c is hydrogen, alkyl, alkenyl, alkynyl, hydroxy, alkoxy, hydroxyalkyl, alkoxyalkyl, aryl, aryloxy, cyano, $-\text{CO}_2\text{R}_a$ or $-\text{C}(\text{O})\text{N}(\text{R}_b)_2$;

Ar is phenyl or naphthyl, optionally and independently mono- or di-substituted with hydroxy, cyano, halogen, alkyl, alkenyl, alkynyl, alkoxy, alkylthio, alkylsulfonyl, arylsulfonyl, amino, nitro, trifluoromethyl, aminomethyl, dimethylaminocarbonyl, benzimidazolyloxy, morpholinocarbonyl, or an aromatic group of formula:



wherein D is oxygen, sulfur, CH_2 , or CO; and each R_d is independently hydrogen, alkyl, alkenyl, alkynyl, alkoxy, trifluoromethyl, amino, halogen, or hydroxy;


R_3 is selected from hydrogen, alkyl, alkenyl, alkynyl, aryl, or heteroaryl; and

R_4 is either a group of formula $-\text{CH}_2\text{-Y}$ or $\text{CHR}_c\text{-Y}$ if R_1 is a group of Formula Set A or a group of formula $\text{CHR}_c\text{-Y}$ if R_1 is a group of Formula Set B;

wherein

R_e is alkyl, alkenyl, alkynyl, hydroxyalkyl, alkoxyalkyl, aminoalkyl, heterocyclalkyl;

Y is a group of formula




wherein R_f is hydrogen, halogen, or alkoxy; and R_g is hydrogen, alkyl, or benzyl.

2. The ligand according to claim 1, wherein R_b is a pyrrolidine, morpholine, piperidine, or piperazine ring, optionally substituted with an alkyl group.
3. The ligand according to claim 1, wherein A and B are oxygen.
4. The ligand according to claim 1, wherein A and B are oxygen; R_1 is a group of Formula Set A wherein four to six atoms separate a basic nitrogen atom from the point of attachment at N3 of the structure of Formula IC; R_2 is a group of formula SO_2 -Ar wherein Ar is an optionally substituted diphenyl ether structure; R_3 is hydrogen; R_e is hydrogen, methyl, or substituted phenyl, Y is an optionally substituted indole ring, R_f is hydrogen, methyl, or halogen, and R_g is hydrogen.
5. The ligand according to claim 1, wherein A and B are oxygen; R_1 is a group of Formula Set B wherein four to six atoms separate a basic nitrogen atom from the point of attachment at N3 of the structure of Formula IC; R_2 is a group of formula SO_2 -Ar wherein Ar is an optionally substituted diphenyl ether structure; R_3 is hydrogen or methyl, R_e is methyl or substituted phenyl, Y is an optionally substituted indole ring, R_f is hydrogen, methyl, or halogen, and R_g is hydrogen.

6. The ligand according to claim 1, wherein structural elements of the R_1 - R_4 groups restrict the conformational mobility of two or more R_1 - R_4 groups appended to the structure of Formula IC.

7. The ligand according to claim 6, wherein two or more of the structural elements are independently: a group of formula $-SO_2-$ attached directly either to the N1 or N3 of the structure of Formula IC; or a branched group having the structure



wherein each R_h independently is any suitable carbon, and the branched group is attached directly to either N1 or C5 of the structure of Formula IC; or a 1,3-cis-disubstituted cyclohexyl group within the R_1 group appended to N3 of the structure of Formula IC.

8. The ligand according to claim 1, wherein the structure of Formula IC contains one or more asymmetric carbon atoms.

9. The ligand according to claim 8, in diastereomeric, racemic, or enantiomeric form.

10. The ligand according to claim 9, having an absolute *S*-stereochemical configuration at C5.

11. The ligand according to claim 1, in free base or pharmaceutically acceptable acid addition salt form.


12. A method for treating a condition in a mammal by the modulation or activation of a somatostatin receptor, the method comprising administering to the mammal an effective amount of a ligand according to claim 1.

13. The method according to claim 12, for treating diabetes and diabetic complications, endocrine disorders, gastrointestinal disorders, CNS disorders, PNS disorders, vascular system disorders, immune system disorders, or cancer.


14. The method according to 13, for treating acromegaly, diarrhea management, diabetic retinopathy, macular degeneration, osteosarcoma, irritable bowel syndrome, or abnormal cellular proliferation.

15. A pharmaceutical composition comprising a compound according to claim 1 and a pharmaceutically acceptable carrier or diluent.

16. A process for the preparation of a compound of Formula IC as defined in claim 1, which includes the steps of reacting a compound of Formula II




wherein A and B are as defined in claim 1 and R'₁, R'₃, and R'₄ are R₁, R₃, and R₄ as defined in claim 1 or protected forms of R₁, R₃, and R₄, with a compound of Formula III



wherein R'₂ is R₂ as defined in claim 1 or is a protected form of R₂ and V is a reactive leaving group, deprotecting the resulting product and recovering the thus obtained compound of Formula IC in free base or acid addition salt form.

17. The process according to claim 16, wherein V is trifluoromethanesulfonate, chlorine, bromine, or iodine.

18. A non-peptide, somatostatin receptor ligand represented by Formula IA:




Formula IA

wherein


A and B independently are O or (H, H);

R₁ is a group of formula



wherein the R_a groups are, independently, hydrogen, alkyl, hydroxyalkyl, alkoxyalkyl, or aryl; X is aryloxy, cyano, -CO₂R_b, or -C(O)N(R_c)₂ wherein R_b is H, alkyl or aryl and wherein the R_c groups are, independently, hydrogen, alkyl, hydroxyalkyl, alkoxyalkyl, aminoalkyl, aryl, heterocycloalkyl or are optionally joined to form a six membered ring; Q is an alkyl spacer group optionally interrupted by oxygen, sulfur or nitrogen atoms and optionally substituted by hydroxy, alkoxy, alkyl, or aryl; Z₁, Z₂, and Z₃ are, independently, carbon, oxygen, sulfur, or nitrogen optionally substituted with an alkyl; n is an integer from 0 to 2; and Y is either hydrogen, an alkyl, heterocycloalkyl, or CH₂N(R_a)₂;

R₂ is -CH(R_d)-Ar or -SO₂-Ar wherein R_d is alkyl, aryl, cyano, -CO₂R_b, or -C(O)N(R_c)₂; Ar is phenyl or naphthyl optionally and independently, mono-




or di-substituted by hydroxy, cyano, halogen, alkyl, alkoxy, -N(R_a)₂, alkylthio, alkylsulfonyl, arylsulfonyl, nitro, -S(O)₂N(R_c)₂, trifluoromethyl, -CH₂N(R_a)₂, -CO₂R_b, -C(O)N(R_c)₂, or an aromatic group of formula

wherein A is CH₂, O, NH, S or CO and the R_e groups, independently, are hydrogen, alkyl, alkoxy, trifluoromethyl, cyano, nitro, amino, halogen, hydroxy;

R₃ is selected from hydrogen, alkyl, aryl, alkylaryl or heteroaryl; and

R₄ is a group of formula:



wherein Z₁, Z₂, and Z₃ are, independently, carbon, oxygen, sulfur or nitrogen optionally substituted with an alkyl; R_f is hydrogen, alkyl or an aryl group; and R_g is hydrogen, halogen, cyano, trifluoromethyl, aryl, alkyl or alkoxy.

19. The ligand according to claim 18, wherein A and B are oxygen.

20. The ligand according to claim 18, wherein R₁ is a group in which 5 or 6 atoms, at least four of which are carbon, separate a basic nitrogen atom from the point of attachment at N3.

21. The ligand according to claim 18, wherein R₂ is a group of formula -SO₂-Ar wherein Ar is an optionally substituted diphenyl ether structure.

22. The ligand according to claim 18, wherein R₃ is hydrogen.

23. The ligand according to claim 18, wherein R₄ is an optionally substituted 3-indolyl group.

24. The ligand according to claim 18, wherein A and B are oxygen, R₁ is a group in which 5 or 6 atoms, at least four of which are carbon, separate a basic nitrogen atom from the

point of attachment at N3, R₂ is -SO₂Ar wherein Ar is an optionally substituted diphenyl ether structure. R₃ is hydrogen, and R₄ is an optionally substituted 3-indolyl group.

25. The ligand according to claim 18, wherein R_c is a morpholine, piperidine, or piperazine ring, optionally substituted with an alkyl group.

26. The ligand according to claim 18, wherein the structure of Formula IA contains one or more asymmetric carbon atoms.

27. The ligand according to claim 26, in diastereomeric, racemic, or enantiomeric form.

28. The ligand according to claim 27, having an absolute *S*-stereochemical configuration at C5.

29. A method for treating a condition in a mammal by the modulation or activation of a somatostatin receptor, the method comprising administering to the mammal an effective amount of a ligand according to claim 18.


30. The method according to claim 29, for treating diabetes and diabetic complications, endocrine disorders, gastrointestinal disorders, CNS disorders, PNS disorders, vascular system disorders, immune system disorders, or cancer.

31. The method according to claim 30, for treating acromegaly, diarrhea management, diabetic retinopathy, macular degeneration, osteosarcoma, irritable bowel syndrome, or abnormal cellular proliferation.

32. A pharmaceutical composition comprising a compound according to claim 18 and a pharmaceutically acceptable carrier or diluent.

33. The liganding according to claim 18, in free base or pharmaceutically acceptable acid addition salt form.


34. The ligand according to claim 18, wherein the ligand has the following structure



wherein

- a is H, OH, NH₂, F, CF₃, Cl, or OCH₃;
- b is H or Cl;
- c is H, C(O)N(CH₃)₂, C(O)N(CH₂CH₂O)₂, CN, or Cl; and
- W is H or CH₃.

35. The ligand according to claim 18, wherein the ligand has the following structure




wherein

- a is OH, NH₂, or F;
- b is H or Cl;

c is H, C(O)N(CH₃)₂, C(O)N(CH₂CH₂O)₂, CN, Cl, or H;
d is CO₂CH₃, CO₂CH₂CH₃, CO₂-tert-butyl, phenyl, CN, and
W is H or CH₃.


36. The ligand according to claim 18, wherein the ligand has the following structure



wherein

a is OH, NH₂, or F;
b is H or Cl; and
c is H, C(O)N(CH₃)₂, C(O)N(CH₂CH₂O)₂, CN, or Cl.

37. A process for the preparation of a compound of Formula IA as defined in claim 18, which includes the steps of reacting a compound of Formula II



wherein A and B are as defined in claim 18 and R₁, R₃, and R₄ are R₁, R₃, and R₄ as defined in claim 18 or protected forms of R₁, R₃, and R₄, with a compound of Formula III


R₂V Formula III

wherein R'₂ is R₂ as defined in claim 18 or is a protected form of R₂ and V is a reactive

leaving group, deprotecting the resulting product and recovering the thus obtained compound of Formula IA in free base or acid addition salt form.

38. The process according to claim 37, wherein V is trifluoromethanesulfonate, chlorine, bromine, or iodine.

39. A non-peptide, somatostatin receptor ligand represented by Formula IB:







Formula IB

wherein


A and B independently are O or (H, H);

Z₁, Z₂, and Z₃ are independently carbon, oxygen, sulfur or a nitrogen atom optionally substituted with an alkyl group;

R₁ is a group of formula




or



wherein X is a saturated or unsaturated aliphatic C₂₋₄ carbonic chain which is (a) optionally interrupted by -O- or -S- or -N(R_b)- and/or (b) optionally substituted by hydroxy, alkoxy groups, alkyl, or aryl; and wherein R_a are independently hydrogen, alkyl, or together form a 3-6 carbon-membered ring; and the R_b independently are hydrogen, alkyl, or arylalkyl; and n is an integer from 0 to 2;

- R₂ is a group of formula -SO₂-Ar or -C(R_c)H-Ar wherein R_c is alkyl, or phenyl optionally substituted by up to three groups alternatively from hydroxy, halogen, alkyl, alkoxy, cyano, trifluoromethyl, amino, monoalkylamino, dialkylamino, carboxy, carboxyalkyl, monoalkylamino-carbonyl, dialkylaminocarbonyl, alkylthio; Ar is phenyl or naphthyl optionally mono- or di-substituted by hydroxy, halogen, alkyl, alkoxy, cyano, trifluoromethyl, aminomethyl, monoalkylaminocarbonyl, dialkylaminocarbonyl, morpholinocarbonyl, piperazinocarbonyl, or a group of formula



wherein A is CH₂, O, S or CO; R_d independently are hydrogen, alkyl, alkoxy, trifluoromethyl, amino, halogen, or hydroxy;

- R₃ is hydrogen, alkyl, aryl, heteroaryl or alkylaryl;
 R₄ is alkyl, aryl, or heteroaryl; and
 R₅ are independently hydrogen, halogen, cyano, alkyl or alkoxyl.

40. The ligand according to claim 39, wherein A and B are oxygen.

41. The ligand according to claim 39, wherein Z₁ and Z₂ are carbon atoms connected by a double bond and Z₃ is preferably N-H.

42. The ligand according to claim 39, wherein R₁ is a group in which 5 atoms separate a basic nitrogen atom from the point of attachment at N3.

43. The ligand according to claim 39, wherein R_2 is a group of formula $-\text{SO}_2\text{-Ar}$ wherein Ar is an optionally substituted diphenyl ether structure.

44. The ligand according to claim 39, wherein R_3 is hydrogen or methyl.

45. The ligand according to claim 39, wherein R_4 is methyl or substituted phenyl.

46. The ligand according to claim 39, wherein R_5 is hydrogen, methyl or halogen.

47. The ligand according to claim 39, wherein A and B are oxygen, R_1 is a group where five atoms separate the point of attachment from a terminal primary amino group, R_2 is a group of formula $-\text{SO}_2\text{-Ar}$ wherein Ar is an optionally substituted diphenyl ether structure, R_3 is hydrogen or methyl, R_4 is methyl or substituted phenyl, and R_5 is hydrogen, methyl or halogen.

48. The ligand according to claim 39, wherein the structure of Formula IB contains one or more asymmetric carbon atoms.

49. The ligand according to claim 48, in diastereomeric, racemic, or enantiomeric form.

50. The ligand according to claim 49, having an absolute *S*-stereochemical configuration at C5.

51. A method for treating a condition in a mammal by the modulation or activation of a somatostatin receptor, the method comprising administering to the mammal an effective amount of a ligand according to claim 39.


52. The method according to claim 51, for treating diabetes and diabetic complications, endocrine disorders, gastrointestinal disorders, CNS disorders, PNS disorders, vascular system disorders, immune system disorders, or cancer.

53. The method according to 52, for treating acromegaly, diarrhea management, diabetic retinopathy, macular degeneration, osteosarcoma, irritable bowel syndrome, or abnormal cellular proliferation.

54. A pharmaceutical composition comprising a compound according to claim 39 and a pharmaceutically acceptable carrier or diluent.

55. The liganding according to claim 39, in free base or pharmaceutically acceptable acid addition salt form.

56. The ligand according to claim 39, wherein the ligand has the following structure



wherein


a is H, OH, NH₂, F, CF₃, Cl, OCH₃, or CF₃;

b is H or Cl;

c is H, C(O)N(CH₃)₂, C(O)N(CH₂CH₂O)₂, CN, or Cl; and

W is H or CH₃.

57. The ligand according to claim 39, wherein the ligand has the following structure




wherein

a is H, NH₂, Cl, CF₃, or OCH₃;

b is H or Cl; and

c is H, C(O)N(CH₃)₂, C(O)N(CH₂CH₂O)₂, CN, and Cl.

58. A process for the preparation of a compound of Formula IB as defined in claim 39, which includes the steps of reacting a compound of Formula II



Formula II

wherein A and B are as defined in claim 39 and R'₁, R'₃, R'₄ and R'₅ are R₁, R₃, R₄ and R₅ as defined in claim 39 or protected forms of R₁, R₃, R₄ and R₅, with a compound of Formula III



wherein R'₂ is R₂ as defined in claim 39 or is a protected form of R₂ and V is a reactive leaving group, deprotecting the resulting product and recovering the thus obtained compound of Formula IB in free base or acid addition salt form.

59. The process according to claim 58, wherein V is trifluoromethanesulfonate, chlorine, bromine, or iodine.

1/2




FIG. 1




FIG. 2

INTERNATIONAL SEARCH REPORT

International Application No.

PCT/US 02/35751

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 A61K31/4166 A61K31/404 A61P35/00 C07D403/06

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D A61P A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, PAJ, BEILSTEIN Data, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 01 85718 A (NOVARTIS ERFINDUNGEN ;BERNEY DANIEL (CH); NEUMANN PETER (CH); NOVA) 15 November 2001 (2001-11-15) claim 1; examples 11-13,16,19-39,45 -----	1-59

☐ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

* Special categories of cited documents:

A document defining the general state of the art which is not considered to be of particular relevance

E earlier document but published on or after the international filing date

L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

O document referring to an oral disclosure, use, exhibition or other means

P document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

& document member of the same patent family

Date of the actual completion of the international search

14 February 2003

Date of mailing of the international search report

26/02/2003

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax (+31-70) 340-3016

Authorized officer

Johnson, C

INTERNATIONAL SEARCH REPORT

Interr. application No.
PCT/US 02/35751

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

Although claims 12-14, 29-31, 51-53 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. ☒ Claims Nos.: 1 (part)
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:

see FURTHER INFORMATION sheet PCT/ISA/210
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
☐ No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Continuation of Box I.2

Claims Nos.: 1 (part)

Present claim 1 relates to compounds defined by reference to a desirable characteristic or property, namely prodrugs. The claims cover all compounds having this characteristic or property, whereas the application provides no support within the meaning of Article 6 PCT and/or disclosure within the meaning of Article 5 PCT for such compounds (the term prodrug does not appear to have been defined at all in the description). In the absence of any structural information to suggest which substituents may be derivatised in what way to give compounds acting as prodrugs, the skilled person would not be able to ascertain the scope of claim 1. Thus claim 1 has only been searched insofar as it relates to compounds of formula IC, their racemates, enantiomers, diastereomers, solvates and salts.

The applicant's attention is drawn to the fact that claims, or parts of claims, relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure.

INTERNATIONAL SEARCH REPORT

Internatio pplication No

PCT/US 02/35751

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 0185718	A	15-11-2001	AU 7399801 A	20-11-2001
			WO 0185718 A1	15-11-2001
			EP 1282617 A1	12-02-2003
<hr/>				